

PATENT APPLICATION TRANSMITTAL LETTER

To the Commissioner of Patents and Trademarks:



CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service as "Express Mail" in an envelope addressed to: Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231, Express Mail Receipt No. EL253288301US

July 1999 Kay Speaker
Date Kay Speaker

Transmitted herewith for filing under 35 U.S.C. 111 and 37 C.F.R §1.53 is the patent application of: Ronald Johnston Hill and Garry Noel Hannan
entitled: "Genetic Sequences Encoding Steroid and Juvenile Hormone Receptor Polypeptides and Insecticidal Modalities Therefor"

Enclosed are:

- ☒ 131 pages of written description, claims, abstract and sequence listing.
- ☒ 4 sheets of drawings.
- ☐ an assignment of the invention to _____
- ☒ unexecuted declaration of the inventor and power of attorney.
- ☐ a certified copy of a _____ application.
- ☐ associate power of attorney.
- ☐ a verified statement to establish small entity status under 37 CFR §1.9 and §1.27.
- ☐ information disclosure statement.
- ☐ preliminary amendment
- ☐ other:

CLAIMS AS FILED

	Number Filed	Number Extra	Rate	Fee
BASIC FEE			\$760	\$760
TOTAL CLAIMS	39-20=	19	x \$18	\$342
INDEPENDENT CLAIMS	13-3=	10	x \$78	\$780
MULTIPLE DEPENDENT CLAIM PRESENT			x \$260	

* Number extra must be zero or larger

TOTAL \$1882

If applicant has small entity status under 37 C.F.R. 1.9 and 1.27, then divide total fee by 2, and enter amount here.

SMALL ENTITY TOTAL

\$

09346470 070199

- ☒ No filing fee is enclosed at this time.
☐ A check in the amount of \$ to cover the filing fee is enclosed.
☐ The Commissioner is hereby authorized to charge and credit Deposit Account No. 07-1969 as described below. A duplicate copy of this sheet is enclosed.

- ☐ Charge the amount of \$_____ as filing fee.
☐ Credit any overpayment.
☐ Charge any additional filing fees required under 37 CFR 1.16 and 1.17.
☐ Charge the issue fee set in 37 CFR 1.18 at the mailing of the Notice of Allowance, pursuant to 37 CFR 1.311(b).
☐ Other _____

☒ Benefit of Prior U.S. Application(s) (35 USC 120)

Applicant claims priority under 35 USC 120 to the following application(s):

PCT application PCT/AU99/00033 filed 15 January 1999


☐ Benefit of Prior U.S. Provisional Application(s) (35 USC 119(e))

Applicant claims priority under 35 USC 120 to the following application(s):

☒ Benefit of Prior Foreign Application(s) (35 USC 119)

Applicant claims priority under 35 USC 119 to the following application:

Australian application PP1356/98 filed 15 January 1998



Donna M. Ferber
Reg. No. 33,878

GREENLEE WINNER and SULLIVAN, P.C.
5370 Manhattan Circle, Suite 201
Boulder, CO 80303
Phone: (303) 499-8080
Fax: (303) 499-8089
email: winner@greenwin.com
ks: July 1, 1999
Docket 53-99

APPLICATION FOR LETTERS PATENT

Inventors:

Ronald Johnston Hill
Garry Noel Hannan

*GENETIC SEQUENCES ENCODING STEROID AND JUVENILE
HORMONE RECEPTOR POLYPEPTIDES AND INSECTICIDAL MODALITIES
THEREFOR*

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service as "Express Mail" in an envelope addressed to: Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231, Express Mail Receipt No. EL253288301US

Date July 1999 Kay Speaker
Kay Speaker

Prepared by:

GREENLEE, WINNER and SULLIVAN, P.C.
5370 Manhattan Circle
Suite 201
Boulder, Colorado 80303
(303) 499-8080

GENETIC SEQUENCES ENCODING STEROID AND JUVENILE HORMONE RECEPTOR POLYPEPTIDES AND INSECTICIDAL MODALITIES THEREFOR

This application is a continuation-in-part of Australian application PP1356/98 filed 15 January
1998 and PCT application PCT/AU99/00033 filed 15 January 1999, both incorporated herein by
reference to the extent that there is no inconsistency with the present disclosure.

FIELD OF THE INVENTION

The present invention relates generally to novel genetic sequences encoding receptor polypeptides
and insecticidal modalities therefor, which insecticidal modalities are based upon non-polypeptide
insect hormones and their receptors. More specifically, the present invention provides isolated
nucleic acid molecules encoding polypeptides comprising functional steroid hormone and juvenile
hormone receptors, in particular isolated nucleic acid molecules which encode polypeptides
comprising the *Lucilia cuprina* and *Myzus persicae* ecdysone receptors and juvenile hormone
receptors. In a particularly preferred embodiment, the present invention relates to isolated nucleic
acid molecules which encode the *L. cuprina* and *M. persicae* EcR polypeptide subunits and EcR
partner protein (USP polypeptide) subunits which form functional heterodimeric ecdysone receptor,
and to the *L. cuprina* and *M. persicae* USP polypeptide of the juvenile hormone receptor. The
present invention further relates to the production of functional recombinant insect receptors and
recombinant polypeptide subunits thereof and derivatives and analogues thereof. The present
invention further relates to the uses of the recombinant receptor and isolated nucleic acid
molecules of the present invention in the regulation of gene expression. The present invention
further relates to screening systems and methods of identifying insecticidally-active agents which
are capable of agonising or antagonising insect receptor function, such as molecules and/or ligands
which associate with steroid receptors or juvenile hormone receptors so as to modify the affinity
of said receptors for their cognate *cis*-acting response elements (e.g., insect steroid response
elements, juvenile hormone response elements) in the genes which they regulate, or alternatively
or in addition, which modify the affinity of said receptors for their cellular stimuli (e.g., insect
steroids or juvenile hormones) or analogues thereof, or alternatively or in addition, which act as
insecticides by virtue of their ability to agonise or antagonise the activity of insect hormones, such
as by mimicry of a ligand which binds to said receptor or a ligand-binding region thereof. The
invention further extends to such compounds and/or ligands.

This specification contains nucleotide and amino acid sequence information prepared using the
program PatentIn Version 2.0, presented herein after the bibliography. Each nucleotide

- 2 -

or amino acid sequence is identified in the sequence listing by the numeric indicator <210> followed by the sequence identifier (e.g. <210>1, <210>2, etc). The length, type of sequence (DNA, protein (PRT), etc) and source organism for each nucleotide or amino acid sequence are indicated by information provided in the numeric indicator fields <211>, <212> and <213>, respectively. Nucleotide and amino acid sequences referred to in the specification are defined by the information provided in numeric indicator field <400> followed by the sequence identifier (eg. <400>1, <400>2, etc).

The designation of nucleotide residues referred to herein are those recommended by the IUPAC-IUB Biochemical Nomenclature Commission, wherein A represents Adenine, C represents Cytosine, G represents Guanine, T represents thymine, Y represents a pyrimidine residue, R represents a purine residue, M represents Adenine or Cytosine, K represents Guanine or Thymine, S represents Guanine or Cytosine, W represents Adenine or Thymine, H represents a nucleotide other than Guanine, B represents a nucleotide other than Adenine, V represents a nucleotide other than Thymine, D represents a nucleotide other than Cytosine and N represents any nucleotide residue.

Bibliographic details of the publications referred to in this specification are collected at the end of the description.

As used herein the term "derived from" shall be taken to indicate that a specified integer may be obtained from a particular source albeit not necessarily directly from that source.

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated step or element or integer or group of steps or elements or integers but not the exclusion of any other step or element or integer or group of elements or integers.

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification,

- 3 -

individually or collectively, and any and all combinations or any two or more of said steps or features.

The present invention is not to be limited in scope by the specific embodiments described
5 herein, which are intended for the purposes of exemplification only. Functionally-equivalent products, compositions and methods are clearly within the scope of the invention, as described herein.

BACKGROUND TO THE INVENTION

10 International Patent Application No WO91/13167 (applicant, The Board of Trustees of Leyland Stanford University, and hereinafter referred to as WO91/13167) describes the identification, characterization, expression and uses of insect steroid receptors and DNA sequences encoding same and, in particular, the identification, characterization, expression and uses of the steroid receptor of the common fruit fly, *Drosophila melanogaster*.

15

It has been found by the present inventors that the limited homology between the *D. melanogaster* steroid receptor-encoding gene sequences and the steroid receptor -encoding sequences derived from other insects, in particular those derived from diptera such as the Australian sheep blowfly *L. cuprina*, hemiptera such as aphids, scale insects and leaf hoppers,
20 coleoptera, neuroptera, lepidoptera, and ants, as well as from helminths and protozoa, prevents the routine isolation of DNA sequences encoding steroid receptors and/or juvenile hormone receptors from these latter-mentioned organisms.

Moreover, the present inventors have discovered that the *D. melanogaster* steroid receptor
25 described in WO91/13167 is temperature-sensitive, showing reduced activity at temperatures above 30°C, such as at temperatures about 37°C, particularly at low concentrations of the receptor. Accordingly, the *D. melanogaster* steroid receptor described in WO91/13167 is of limited utility at physiological temperatures applicable to animal or bacterial cells. Moreover, wherein it is desirable to produce a biologically-active steroid receptor using *in vivo* or *in situ*
30 expression systems, which expression systems routinely utilise cells or tissues in the temperature range of about 28°C to about 42°C, the *D. melanogaster* steroid receptor is also

- 4 -

of limited utility.

In work leading up to the present invention, the present inventors developed a novel screening protocol, involving the utilisation of highly-degenerate oligonucleotide probes and primers
5 derived from the amino acid sequences of the DNA-binding domains of the *D. melanogaster* and *Chironomus tentans* ecdysone receptor polypeptides, to identify nucleotide sequences encoding novel steroid receptor polypeptides and novel insect juvenile hormone receptor polypeptides. The present inventors have further identified specific regions within these novel polypeptides which are suitable for use in preparing a surprising range of novel steroid
10 receptor polypeptide derivatives and insect juvenile hormone receptor polypeptide derivatives. The novel steroid receptor polypeptides and novel insect juvenile hormone receptor polypeptides of the present invention, and derivative polypeptides thereof, and assembled steroid receptors and insect juvenile hormone receptors derived from said polypeptides and derivatives, and nucleic acid molecules encoding same as exemplified herein, provide the
15 means for developing a wide range of insecticidally-active agents, as well as methods for the regulated production of bioactive molecules. In particular, the present invention provides the means for developing specific ligands which bind to and either agonise or antagonise the steroid receptors and/or juvenile hormone receptors and/or polypeptide subunits thereof as described herein, thereby functioning as highly-specific insecticides, offering significant
20 commercial and environmental benefits.

The present inventors have been surprisingly successful in characterizing the ecdysone receptor and juvenile hormone receptor derived from insects of the orders Diptera and Hemiptera, and polypeptide components thereof and functional derivatives of said polypeptides
25 and receptors, particularly in light of the extreme difficulties in dealing with these organisms. The nature of these molecules was unknown prior to the present invention.

The various aspects of this invention overcome the problems associated with *Drosophila* ecdysone receptors which lack thermal stability. Moreover, those aspects of the invention
30 pertaining to methods of screening for insecticidally active agents do not involve competition assays which are generally complex, and often inaccurate or difficult to calibrate.

- 5 -

SUMMARY OF THE INVENTION

One aspect of the present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence which encodes or is complementary to a sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide or a bioactive derivative or analogue thereof, wherein said polypeptide:

- (i) is selected from the list comprising EcR polypeptide of an steroid receptor, the partner protein (USP polypeptide) of an steroid receptor and the USP polypeptide of a juvenile hormone receptor; and
- (ii) comprises an amino acid sequence that is at least 40% identical to any one of the amino acid sequences set forth in <400>2, <400>4, <400>6, <400>10, <400>12 or <400>14.

In an alternative embodiment, the present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence which encodes or is complementary to a sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide or a bioactive derivative or analogue thereof, wherein said polypeptide:

- (i) is selected from the list comprising EcR polypeptide of an steroid receptor, the partner protein (USP polypeptide) of an steroid receptor and the USP polypeptide of a juvenile hormone receptor; and
- (ii) comprises an amino acid sequence that is at least 40% identical to any one of the amino acid sequences encoded by the plasmids deposited under any one of AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568.

In a further alternative embodiment, the isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which encodes or is complementary to a sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide or a bioactive derivative or analogue thereof, wherein said nucleotide sequence is selected from the list comprising:

- (i) a nucleotide sequence having at least 40% identity to any one of the nucleotide sequences set forth in <400>1, <400>3, <400>5, <400>9, <400>11 or <400>13 or a complementary nucleotide sequence thereto;

- 6 -

5 (iii) a nucleotide sequence that is capable of hybridising under at least low stringency conditions to a nucleotide sequence contained in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568.;

(v) a nucleotide sequence that is amplifiable by PCR using a nucleic acid primer sequence set forth in any one of <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20.

20 In a further alternative embodiment, the present invention provides an isolated nucleic acid molecule which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide and comprises the nucleotide sequence set forth in <400>3 or <400>13 or a complementary nucleotide sequence thereto.

25 In a further alternative embodiment, the present invention provides an isolated nucleic acid molecule which encodes a steroid receptor polypeptide and comprises the nucleotide sequence set forth in <400>5 or <400>7 or <400>8 or <400>9 or a complementary nucleotide sequence thereto.

30 In a further alternative embodiment, the present invention provides an isolated nucleic acid molecule which encodes a steroid receptor polypeptide or a juvenile hormone receptor

- 7 -

polypeptide and comprises the nucleotide sequence set forth in <400>11 or a complementary nucleotide sequence thereto.

A second aspect of the present invention provides a method of identifying an isolated nucleic acid molecule which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide comprising the steps of:

(i) hybridising genomic DNA, mRNA or cDNA with a hybridisation-effective amount of one or more probes selected from the list comprising:

(a) probes comprising at least 10 contiguous nucleotides in length derived from any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, <400>13, <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20 or a complementary nucleotide sequence thereto;

(b) probes comprising at least 10 contiguous nucleotides in length derived from a cDNA contained in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568; and

(c) hybridisation probes comprising the nucleotide sequences set forth in any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, or <400>13 or a complementary nucleotide sequence thereto or a homologue, analogue or derivative thereof which is at least 40% identical to said sequence or complement; and

(ii) detecting the hybridisation.

In an alternative embodiment, the inventive method comprises the steps of:

(i) annealing to genomic DNA, mRNA or cDNA, one or more PCR primers comprising at least 10 contiguous nucleotides in length derived from the group consisting of:

(a) a primer derived from any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, <400>13, <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20 or a complementary nucleotide sequence thereto; and

(b) a primer derived from a cDNA contained in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566,

667620 "04434660

- 8 -

NM99/04567, or NM99/04568; and

- (ii) amplifying a nucleotide sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide in a polymerase chain reaction.

5 In a further alternative embodiment, the inventive method comprises the steps of:

- (i) amplifying a nucleotide sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide in a polymerase chain reaction using one or more PCR primers comprising at least 10 contiguous nucleotides in length from the group consisting of:

- 10 (a) a primer derived from any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, <400>13, <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20 or a complementary nucleotide sequence thereto; and
 (b) a primer derived from a cDNA contained in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566,
 15 NM99/04567, or NM99/04568;

- (ii) hybridising the amplified nucleotide sequence to genomic DNA, mRNA or cDNA with a hybridisation-effective amount of one or more probes selected from the group consisting of:

- 20 (a) a probe comprising at least 10 contiguous nucleotides in length derived from any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, <400>13, <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20 or a complementary nucleotide sequence thereto;
 (b) a probe comprising at least 10 contiguous nucleotides in length derived from a cDNA contained in any one of the plasmids deposited under AGAL
 25 Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568; and
 (c) hybridisation probes comprising the nucleotide sequences set forth in any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, or <400>13 or a complementary nucleotide sequence thereto or a homologue, analogue or derivative thereof which is at least 40% identical to said sequence
 30 or complement; and
 (iii) detecting the hybridisation.

664620 "0434630

- 9 -

A third aspect of the present invention provides a genetic construct comprising the subject isolated nucleic acid molecule which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide, operably linked to a promoter sequence. Preferably, the subject
5 nucleic acid molecule is in an expressible format, such that it is possible to produce a recombinant polypeptide therefrom.

Accordingly, a fourth aspect of the invention provides a recombinant or isolated polypeptide comprising a steroid receptor polypeptide or juvenile hormone receptor polypeptide or a
10 bioactive derivative or analogue thereof, wherein said polypeptide:

- (i) is selected from the list comprising EcR polypeptide of a steroid receptor, the partner protein (USP polypeptide) of a steroid receptor and the USP polypeptide of a juvenile hormone receptor; and
- (ii) comprises an amino acid sequence that is at least 40% identical to any one of
15 the amino acid sequences set forth in <400>2, <400>4, <400>6, <400>10, <400>12 or <400>14;

wherein said polypeptide is substantially free of naturally-associated insect cell components.

In an alternative embodiment, the invention provides a recombinant or isolated polypeptide
20 comprising a steroid receptor polypeptide or juvenile hormone receptor polypeptide or a bioactive derivative or analogue thereof, wherein said polypeptide:

- (i) is selected from the list comprising EcR polypeptide of a steroid receptor, the partner protein (USP polypeptide) of a steroid receptor and the USP polypeptide of a juvenile hormone receptor; and
- (ii) comprises an amino acid sequence that is at least 40% identical to a
25 polypeptide encoded by the cDNA present in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568;

wherein said polypeptide is substantially free of naturally-associated insect cell components.

30 A fifth aspect of the invention provides a cell comprising the subject isolated nucleic acid molecule which encodes a steroid receptor polypeptide or a juvenile hormone receptor

- 10 -

polypeptide.

In a preferred embodiment, the cell of the present invention expresses the polypeptide encoded by the nucleic acid molecule.

5

In a preferred embodiment, the cell expresses a steroid receptor polypeptide or a fragment thereof which receptor is capable of binding to an insect steroid or analogue thereof or a candidate insecticidally active agent to form an activated complex, and comprises a nucleic acid sequence encoding a bioactive molecule or a reporter molecule operably linked to one
10 or more insect steroid response elements which on binding of the said activated complex promotes transcription of the nucleic acid sequence, wherein said cell on exposure to insect steroid or an analogue thereof, regulates expression of said bioactive molecule or allows detection of said reporter molecule.

15 In a further aspect of this invention, there is provided an animal (such as a mammal), microorganism, plant or aquatic organism, containing one or more cells as mentioned above.

A further aspect of the present invention provides a method of identifying a modulator of insect steroid receptor-mediated gene expression or insect juvenile hormone receptor-mediated gene
20 expression comprising:

- (i) assaying the expression of a reporter gene in the presence of a recombinant or isolated insect steroid receptor polypeptide or a juvenile hormone receptor polypeptide of the invention and a potential modulator; and
 - (ii) assaying the expression of a reporter gene in the presence of a recombinant
25 or isolated insect steroid receptor polypeptide or a juvenile hormone receptor polypeptide of the invention and without said potential modulator; and
 - (ii) comparing expression of the reporter gene in the presence of the potential modulator to the expression of a reporter gene in the absence of the potential modulator,
- 30 wherein said reporter gene is placed operably under the control of a steroid response element (SRE) to which said insect steroid receptor binds or a promoter sequence comprising said

- 11 -

SRE.

A still further aspect of the invention provides a method of identifying a potential insecticidal compound comprising:

- 5 (i) assaying the binding directly or indirectly of a recombinant or isolated insect steroid receptor polypeptide or a juvenile hormone receptor polypeptide of the invention to a steroid response element (SRE) to which said insect steroid receptor binds, in the presence of a candidate compound; and
- 10 (ii) assaying the binding directly or indirectly of a recombinant or isolated insect steroid receptor polypeptide or a juvenile hormone receptor polypeptide of the invention to a steroid response element (SRE) to which said insect steroid receptor binds, in the absence of said candidate compound; and
- 15 (ii) comparing the binding assayed at (i) and (ii), wherein a difference in the level of binding indicates that the candidate compound possesses potential insecticidal activity.

A still further aspect of the invention provides a method of identifying a candidate insecticidally-active agent comprising the steps of:

- 20 a) expressing an EcR polypeptide of a steroid receptor or a fragment thereof which includes the ligand-binding region, optionally in association with an EcR partner protein (USP polypeptide) of a steroid receptor or ligand binding domain thereof, optionally in association with an insect steroid or analogue thereof so as to form a complex;
- 25 b) purifying or precipitating the complex;
- c) determining the three-dimensional structure of the ligand binding domain of the complex; and
- d) identifying compounds which bind to or associate with the three-dimensional structure of the ligand binding domain, wherein said compounds represent candidate insecticidally-active agents.
- 30

A still further aspect of the invention provides a method of identifying a candidate insecticidally-

- 12 -

active agent comprising the steps of:

- a) expressing a USP polypeptide of a juvenile hormone receptor or a fragment thereof which includes the ligand-binding region, optionally in association with an EcR polypeptide of a steroid receptor or ligand binding domain thereof, and optionally in association with an insect steroid or analogue thereof, so as to form a complex;
- b) purifying or precipitating the complex;
- c) determining the three-dimensional structure of the ligand binding domain of the complex; and
- d) identifying compounds which bind to or associate with the three-dimensional structure of the ligand binding domain, wherein said compounds represent candidate insecticidally-active agents.

In another aspect this invention relates to a method or assay for screening insecticidally active compounds which comprises reacting a candidate insecticidal compound with a steroid receptor polypeptide or fragment thereof encompassing the ligand binding domain, or complex thereof with a partner protein or a fragment thereof which encompasses the ligand binding domain, and detecting binding or absence of binding of said compound so as to determine insecticidal activity.

- A still further aspect of the invention provides a synthetic compound which interacts with the three dimensional structure of a polypeptide or protein selected from the list comprising:
- (i) an EcR polypeptide of a steroid receptor or a fragment thereof;
 - (ii) an EcR partner protein (USP polypeptide) of a steroid receptor or a fragment thereof;
 - (iii) a USP polypeptide of a juvenile hormone receptor; and
 - (iv) a functional receptor or protein complex formed by association of (i) and (ii),
- wherein said compound is capable of binding to said polypeptide or protein to agonise or antagonise the binding activity or bioactivity thereof.
- Preferably, the synthetic compounds are derived from the three dimensional structure of insect steroid receptor(s) or juvenile hormone receptor(s) which compounds bind to said receptor(s)

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 13 -

and have the effect of either inactivating the receptor(s) or potentiating the activity of the receptor(s). More preferably, the compounds mimic the three-dimensional structure of a ligand which binds to the receptor(s) and more preferably, mimic the three-dimensional structure of a ligand which binds to the ligand-binding region of said receptor(s).

5

In a still further aspect of this invention, there is provided a screening system for insecticidally active agents comprising a nucleotide sequence encoding a steroid receptor or a fragment thereof, and a nucleotide sequence encoding a partner protein or a fragment thereof which associates with the receptor so as to confer enhanced affinity for insect steroid response
10 elements, enhanced affinity for insect steroids or analogues thereof, or insecticidally active agents and/or thermostability or enhanced thermostability of said receptor, which receptor and partner protein is capable of binding to a candidate insecticidally active agent to form an activated complex, and a nucleic acid sequence encoding a bioactive molecule or a reporter molecule operably linked to one or more insect steroid response elements which on binding
15 of the said activated complex regulates transcription of the nucleic acid sequence, wherein on exposure to said agent expression of the bioactive molecule or reporter molecule correlates with insecticidal activity.

In another aspect of this invention, there is provided a method for the regulated production of
20 a bioactive molecule or a reporter molecule in a cell, said method comprising the steps of introducing into said cell:

- a) a nucleotide sequence encoding a steroid receptor or a fragment thereof which is capable of binding an insect steroid or analogue thereof, to form an activated complex; and
- 25 b) a nucleotide sequence encoding said bioactive molecule or reporter molecule operably linked to one or more insect steroid response elements which on binding of the said activated complex regulates transcription of the nucleic acid sequence encoding said bioactive molecule or reporter molecule,

wherein exposing the cell to an insect steroid or analogue thereof regulates expression of the
30 bioactive molecule or reporter molecule.

- 14 -

SUMMARY OF SEQUENCE LISTING

- <400>1:** The nucleotide sequence of the open reading frame of a cDNA molecule which encodes the EcR polypeptide subunit of the *L. cuprina* ecdysone receptor and amino acid sequence therefor.
- 5 **<400>2:** The amino acid sequence of the EcR polypeptide subunit of the *L. cuprina* ecdysone receptor.
- <400>3:** The nucleotide sequence of the open reading frame of a cDNA molecule which encodes the EcR partner protein (USP polypeptide) subunit of the *L. cuprina* ecdysone receptor and/or which encodes the USP polypeptide subunit of the
- 10 **<400>4:** The amino acid sequence of the EcR partner protein (USP polypeptide) subunit of the *L. cuprina* ecdysone receptor and/or the amino acid sequence of the USP polypeptide subunit of the *L. cuprina* juvenile hormone receptor.
- <400>5:** The nucleotide sequence of a cDNA molecule which encodes part of the EcR polypeptide subunit of the *M. persicae* ecdysone receptor and amino acid
- 15 **<400>6:** The amino acid sequence of a part of the EcR polypeptide subunit of the *M. persicae* ecdysone receptor.
- <400>7:** The nucleotide sequence of the EcR probe 1 which is specific for genetic sequences encoding the EcR polypeptide subunit of aphid ecdysone receptors, in particular the EcR polypeptide subunit of the *M. persicae* ecdysone receptor.
- 20 **<400>8:** The nucleotide sequence of the EcR probe 2 sequence which is specific for genetic sequences encoding the EcR polypeptide subunit of aphid ecdysone receptors, in particular the EcR polypeptide subunit of the *M. persicae* ecdysone receptor.
- 25 **<400>9:** The nucleotide sequence of the open reading frame of a cDNA molecule which encodes the EcR polypeptide subunit of the *M. persicae* ecdysone receptor and amino acid sequence therefor.
- <400>10:** The amino acid sequence of the EcR polypeptide subunit of the *M. persicae* ecdysone receptor.
- 30 **<400>11:** The nucleotide sequence of the open reading frame of a cDNA molecule which

65402002131650

- 15 -

encodes the EcR partner protein (USP polypeptide) subunit of the *M. persicae* ecdysone receptor and/or which encodes the USP polypeptide subunit of the *M. persicae* juvenile hormone receptor, and amino acid sequence therefor.

5 **<400>12:** The amino acid sequence of the EcR partner protein (USP polypeptide) subunit of the *M. persicae* ecdysone receptor and/or the amino acid sequence of the USP polypeptide subunit of the *M. persicae* juvenile hormone receptor.

10 **<400>13:** The nucleotide sequence of a 150 base-pair probe which is specific for genetic sequences encoding the EcR partner protein (USP polypeptide) subunit of *L. cuprina* ecdysone receptor and/or the USP polypeptide subunit of the *L. cuprina* juvenile hormone receptor, and amino acid sequence therefor.

15 **<400>14:** The amino acid sequence encoded by the nucleotide sequence of <400>13, comprising amino acid residues 108-149 of the EcR partner protein (USP polypeptide) subunit of the *L. cuprina* ecdysone receptor and/or amino acid residues 108-149 of the amino acid sequence of the USP polypeptide subunit of the *L. cuprina* juvenile hormone receptor set forth herein as <400>4.

<400>15: The nucleotide sequence of the degenerate primer Rdna3.

<400>16: The nucleotide sequence of the degenerate primer Rdna4.

<400>17: The nucleotide sequence of the primer Mdna1.

<400>18: The nucleotide sequence of the primer Mdna2.

20 **<400>19:** The nucleotide sequence of the primer AP1.

<400>20: The nucleotide sequence of the degenerate primer AP2.

BRIEF DESCRIPTION OF THE DRAWINGS

25 **Figure 1** is a graphical representation showing function of the EcR polypeptide subunit of the *L. cuprina* ecdysone receptor *in vivo*. CHO cells were cotransfected with:

- (1) one of the following expression plasmids: pSGDmEcR, pSGLcEcR, or the parental expression plasmid pSG5 as a control, at 1 µg/ml;
- (2) plasmid p(EcRE)₇-CAT (1 µg/ml); and
- 30 (3) an independent reporter plasmid, pPGKLacZ, at 1 µg/ml.

CAT expression was induced with Muristerone A at either 10 µM or 50 µM while control cells

034440 "02434800
667020 "02434800

- 16 -

received only the carrier ethanol. ELISA kits were used to quantify the synthesis of CAT and β -galactosidase in extracts of cells forty eight hours after transfection. The level of CAT was normalized to the level of β -galactosidase in the same extract. Fold-induction represents the normalized values for CAT gene expression in cells transfected with pSGDmEcR, pSGLcEcR or pSG5 in the presence of hormone, relative to the normalized values for CAT gene expression in cells transfected with the same plasmid, but in the absence of hormone. The average values of three independent experiments are shown and the error bars indicate standard error of the mean.

- 10 **Figure 2** is a copy of a graphical representation showing the activity of plasmid pSGLD and pSGDL, containing chimeric EcR polypeptide subunits of insect ecdysone receptors, produced as described in the Examples. Cotransfection assays were performed as described in the Examples using plasmids pSGLD and pSGDL and the CAT reporter plasmid p(EcRE)₇-CAT (1ug/ml) and an independent reporter, pPGKLacZ at 1 μ g/ml each. CAT/b-Gal (%) refers to CAT reporter activity expressed as a percentage relative to β -galactosidase activity produced by the internal control reporter, pPGKLacZ.

Figure 3 is a copy of a graphical representation showing the binding activity in extracts of Sf9 and Sf21 cells containing a baculovirus expressing LcEcRDEF and LcUSPDEF, as described in the Examples. Control cells contained baculovirus expressing β -galactosidase and CAT only.

Figure 4 is a graphical representation showing the ecdysteroid binding activities of an *in vitro*-translated *Myzus persicae* EcR (MpEcR) polypeptide, an *in vitro*-translated *Myzus persicae* USP (MpUSP) polypeptide, and an *in vitro*-translated complex of the *M.persicae* EcR and USP polypeptides.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

One aspect of the present invention provides an isolated nucleic acid molecule comprising a nucleotide sequence which encodes or is complementary to a sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide or a bioactive

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 17 -

derivative or analogue thereof, wherein said polypeptide:

- (i) is selected from the list comprising EcR polypeptide of a steroid receptor, the partner protein (USP polypeptide) of a steroid receptor and the USP polypeptide of a juvenile hormone receptor; and
- 5 (ii) comprises an amino acid sequence that is at least 40% identical to any one of the amino acid sequences set forth in <400>2, <400>4, <400>6, <400>10, <400>12 or <400>14.

Accordingly, the isolated nucleic acid molecule of the invention may comprise a fragment of
10 a nucleotide sequence encoding a full-length receptor polypeptide.

It is to be understood that a "fragment" of a nucleotide sequence encoding an EcR polypeptide subunit of a steroid receptor or an EcR partner protein (USP polypeptide) of a steroid receptor or a USP polypeptide of a juvenile hormone receptor, refers to a nucleotide sequence
15 encoding a part or fragment of such a receptor which is capable of binding or associating with an insect steroid or an analogue thereof, or a candidate insecticidally active compound. Fragments of a nucleotide sequence would generally comprise in excess of twenty contiguous nucleotides derived from the base sequence and may encode one or more domains of a functional insect steroid receptor or juvenile hormone receptor.

20

Preferably, the isolated nucleic acid molecule of the invention encodes an ecdysteroid receptor polypeptide. Those skilled in the art are aware that ecdysteroid receptors derived from insects are heterodimeric receptors comprising an EcR polypeptide subunit and an EcR partner protein (USP polypeptide) (see also Jones and Sharp, 1997). In this regard, the present
25 inventors have discovered that the USP polypeptide of the insect juvenile hormone receptor is structurally-identical to the EcR partner protein of the ecdysteroid receptor of the present invention, however juvenile hormone receptors comprise monomers or multimers of the USP polypeptide acting without the EcR polypeptide subunit that is present in the ecdysteroid receptors. Accordingly, the present invention extends equally to nucleotide sequences
30 encoding polypeptides of both the ecdysteroid receptors and polypeptides of the juvenile hormone receptors of insects.

- 18 -

More preferably, the isolated nucleic acid molecule of the invention encodes an ecdysteroid receptor that is modulated by one or more of the steroids ecdysone, ponasterone A, or muristerone, or an analogue of an ecdysteroid.

- 5 The isolated nucleic acid molecule of the invention may be derived from any organism that contains steroid receptors that are responsive to ecdysteroids or ecdysteroid-like compounds or juvenile hormones, or analogues of such receptor-ligands. Accordingly, the present invention is not to be limited in any of its embodiments to the particular source of the subject nucleic acid, or polypeptide encoded therefor.

10

Preferably, the isolated nucleic acid molecule of the invention is derived from insects, helminths (nematodes, cestodes, trematodes), protozoa, and ants, amongst others.

More preferably, the isolated nucleic acid molecule of the invention is derived from an insect

- 15 selected from the list comprising diptera, hemiptera, coleoptera, neuroptera, lepidoptera and ants, amongst others. Still more preferably, the isolated nucleic acid molecule of the present invention is derived from aphids, scale insects, leaf hoppers, white fly, and blowflies such as sheep blowflies.

- 20 The present invention does not extend to amino acid sequences comprising the complete EcR polypeptide subunit of the *D. melanogaster* ecdysone receptor as described in WO91/13167. However, this exclusion is made on the understanding that the present invention does encompass chimeric genes and fusion proteins which include the *D. melanogaster* nucleotide and amino acid sequences, respectively.

25

In a particularly preferred embodiment, the isolated nucleic acid molecule of the present invention is derived from the aphid *M. persicae* or alternatively, from the Australian sheep blowfly, *L. cuprina*.

- 30 The ecdysteroid receptor is preferably modulated by one or more of the steroids ecdysone, ponasterone A, or muristerone, or an analogue of an ecdysteroid.

- 19 -

As used herein, the term "analogue of an ecdysteroid" shall be taken to indicate any compound that binds to one or more polypeptide subunits of an ecdysteroid receptor or the heterodimeric holoreceptor comprising same or alternatively or in addition, which binds to the USP polypeptide of a juvenile hormone receptor or alternatively or in addition, which binds to a bioactive derivative or analogue of said polypeptides or holoreceptor. The term "analogue of an ecdysteroid" shall further be taken to indicate any compound that modulates the bioactivity of one or more polypeptide subunits of an ecdysteroid receptor or the heterodimeric holoreceptor comprising same or alternatively or in addition, that modulates the bioactivity of the USP polypeptide of a juvenile hormone receptor or alternatively or in addition, that modulates the bioactivity of a bioactive derivative or analogue of said polypeptides or holoreceptor.

The present invention is not to be limited in scope to the specific *L. cuprina* and *M. persicae* nucleotide and amino acid sequences set forth in the accompanying Sequence Listing and persons skilled in the art will readily be able to identify additional related sequences from other sources using art-recognised procedures, for example using nucleic acid hybridisation and/or polymerase chain reaction essentially as described by Ausubel *et al.* (1992) and/or McPherson *et al.* (1991) and/or Sambrook *et al.* (1989).

Accordingly, the present invention clearly encompasses isolated nucleic acid molecules which encode or are complementary to isolated nucleic acid molecules which encode the subject EcR polypeptide of a steroid receptor or fragments thereof, and/or the subject EcR partner proteins (USP polypeptide) of a steroid receptor and/or the subject USP polypeptide of a juvenile hormone receptor, in addition to derivatives, fragments and analogues thereof which comprise amino acid sequences having at least 40% identity to the amino acid sequences set forth in any one of <400>2, <400>4, <400>6, <400>10, <400>12 or <400>14.

The present invention clearly extends further to isolated nucleic acid molecules which encode or are complementary to isolated nucleic acid molecules which encode the subject EcR polypeptide of a steroid receptor or fragments thereof, and/or the subject EcR partner proteins (USP polypeptide) of a steroid receptor and/or the subject USP polypeptide of a juvenile

- 20 -

hormone receptor, in addition to derivatives, fragments and analogues thereof which comprise amino acid sequences having at least 40% identity to any one or more of the amino acid sequences encoded by the *L. cuprina* or *M. persicae* cDNAs contained in AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568.

5

For the purposes of nomenclature, plasmid pLcEcR contains the cDNA encoding the EcR polypeptide subunit of the *Lucillia cuprina* ecdysone receptor. This plasmid was deposited on 1 July, 1999 with the Australian Government Analytical Laboratories at 1 Suakin Street, Pymble, New South Wales 2073, Australia under the provisions of the Budapest Treaty on the
10 International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and accorded AGAL Accession No. NM99/04566.

For the purposes of nomenclature, plasmid pLcUSP contains the cDNA encoding the EcR partner protein (USP polypeptide) subunit of the *Lucillia cuprina* ecdysone receptor and/or the
15 USP polypeptide subunit of the *L. cuprina* juvenile hormone receptor. This plasmid was deposited on 1 July, 1999 with the Australian Government Analytical Laboratories at 1 Suakin Street, Pymble, New South Wales 2073, Australia under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and accorded AGAL Accession No. NM99/04565.

20

For the purposes of nomenclature, plasmid pMpEcR contains the cDNA encoding the EcR polypeptide subunit of the *Myzus persicae* ecdysone receptor. This plasmid was deposited on 1 July, 1999 with the Australian Government Analytical Laboratories at 1 Suakin Street, Pymble, New South Wales 2073, Australia under the provisions of the Budapest Treaty on the
25 International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and accorded AGAL Accession No. NM99/04567.

For the purposes of nomenclature, plasmid pMpUSP contains the cDNA encoding the EcR partner protein (USP polypeptide) subunit of the *Myzus persicae* ecdysone receptor and/or the
30 USP polypeptide subunit of the *M. persicae* juvenile hormone receptor. This plasmid was deposited on 1 July, 1999 with the Australian Government Analytical Laboratories at 1 Suakin

667040 "0249660

- 21 -

Street, Pymble, New South Wales 2073, Australia under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure and accorded AGAL Accession No. NM99/04568.

- 5 The deposits referred to herein will be maintained under the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. These deposits are provided merely for the purposes of exemplification and are not an admission that a deposit is required under 35USC §112. A license may be required to make, use or sell the deposited materials or a polypeptide encoded by a cDNA thereof and no
- 10 such license is hereby granted. It is to be understood however, that the deposits will become publicly available upon the grant of a patent pertaining to the instant disclosure in so far as that patent relates to the deposits referred to herein.

- Preferably, the percentage similarity to any one of <400>2, <400>4, <400>6, <400>10,
- 15 <400>12 or <400>14, or to a polypeptide encoded by a cDNA contained in any one of AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568 is at least about 60%, more preferably at least about 80%, even more preferably at least about 90%.

- In determining whether or not two amino acid sequences fall within these percentage limits,
- 20 those skilled in the art will be aware that it is necessary to conduct a side-by-side comparison or multiple alignment of sequences. In such comparisons or alignments, differences will arise in the positioning of non-identical residues, depending upon the algorithm used to perform the alignment. In the present context, reference to a percentage identity or similarity between two or more amino acid sequences shall be taken to refer to the number of identical and similar
- 25 residues respectively, between said sequences as determined using any standard algorithm known to those skilled in the art. For example, amino acid sequence identities or similarities may be calculated using the GAP programme and/or aligned using the PILEUP programme of the Computer Genetics Group, Inc., University Research Park, Madison, Wisconsin, United States of America (Devereaux *et al*, 1984). The GAP programme utilizes the algorithm of
- 30 Needleman and Wunsch (1970) to maximise the number of identical/similar residues and to minimise the number and/or length of sequence gaps in the alignment. Alternatively or in

- 22 -

addition, wherein more than two amino acid sequences are being compared, the ClustalW programme of Thompson *et al* (1994) is used.

In an alternative embodiment, the isolated nucleic acid molecule of the invention encodes or
5 is complementary to an isolated nucleic acid molecule which encodes a steroid receptor polypeptide or a fragment thereof, or a partner protein (USP) or a fragment thereof, which at least comprises an amino acid sequence which is substantially identical to any one of <400>2, <400>4, <400>6, <400>10, <400>12 or <400>14 or to the amino acid sequence of a polypeptide encoded by any one of the cDNAs contained in AGAL Accession Nos.
10 NM99/04565, NM99/04566, NM99/04567, or NM99/04568.

As used herein, the term "substantially identical" or similar term shall be taken to include any sequence which is at least about 95% identical and preferably at least 99% or 100% identical to a stated nucleotide sequence or amino acid sequence, including any homologue, analogue
15 or derivative of said stated nucleotide sequence or amino acid sequence.

Those skilled in the art will be aware that variants of the nucleotide sequences set forth in any one of <400>1 or <400>3 or <400>5 or <400>7 or <400>8 or <400>9 or <400>11 or <400>13, or variants of the cDNAs contained in any one of the deposited plasmids, which variants
20 encode EcR polypeptides of insect steroid receptors or fragments thereof or EcR partner proteins (USP polypeptides) or fragments thereof, or USP polypeptides of insect juvenile hormone receptors, may be isolated by hybridization under low stringency conditions as exemplified herein.

25 Such variants include any genomic sequences, cDNA sequences mRNA or other isolated nucleic acid molecules derived from the nucleic acid molecules exemplified herein by the Sequence Listing. Additional variants are not excluded.

In a particularly preferred embodiment of the invention, the variant nucleotide sequences
30 encode a fragment of the EcR polypeptide of the insect steroid receptor or a fragment of the EcR partner protein (USP polypeptide) of the insect steroid receptor or a fragment of the USP

P:\OPER\IMRO\ECDYSONE.CIP - 1/7/99

- 23 -

polypeptide of the insect juvenile hormone receptor.

Preferred fragments of the subject polypeptides include one or more regions or domains which are involved in the interaction or association between the monomeric polypeptide subunits of a multimeric receptor and/or which are involved in the interaction or association between (i) a cognate steroid or receptor ligand or *cis*-acting DNA sequence; and (ii) said monomeric polypeptide subunits or the receptor *per se*. In a particularly preferred embodiment, the fragments comprise the DNA-binding domain, linker domain (domain D) or a part thereof, or ligand-binding domain (eg. hormone-binding domain) of a steroid receptor polypeptide or juvenile hormone receptor polypeptide or receptor holoenzyme. As exemplified herein, wherein biological activity of the *L. cuprina* ecdysone receptor is required, it is preferably to include at least a ligand-binding region comprising the ligand-binding domain and at least a part of the linker domain of the EcR polypeptide subunit, optionally in association with a ligand-binding region comprising at least the ligand-binding domain and at least a part of the linker domain of the EcR partner protein (USP polypeptide) subunit of said receptor. Additional fragments are not excluded.

Homologues, analogues and derivatives of the nucleotide sequences exemplified herein may be isolated by hybridising same under at least low stringency conditions and preferably under at least medium stringency conditions, to the nucleic acid molecule set forth in any one of <400>1 or <400>3 or <400>5 or <400>7 or <400>8 or <400>9 or <400>11 or <400>13 or to a complementary strand thereof, or to the cDNAs contained in any one or more of the deposited plasmids. More preferably, the isolated nucleic acid molecule according to this aspect of the invention is capable of hybridising under at least high stringency conditions to the nucleic acid molecule set forth in any one of <400>1 or <400>3 or <400>5 or <400>7 or <400>8 or <400>9 or <400>11 or <400>13 or to a complementary strand thereof, or to the cDNAs contained in any one or more of the deposited plasmids.

For the purposes of defining the level of stringency, a low stringency is defined herein as being a hybridisation and/or a wash carried out in 6xSSC buffer, 0.1% (w/v) SDS at 28°C or alternatively, as exemplified herein. Generally, the stringency is increased by reducing the

- 24 -

concentration of SSC buffer, and/or increasing the concentration of SDS and/or increasing the temperature of the hybridisation and/or wash. A medium stringency comprises a hybridisation and/or a wash carried out in 0.2xSSC-2xSSC buffer, 0.1% (w/v) SDS at 42°C to 65°C, while a high stringency comprises a hybridisation and/or a wash carried out in 0.1xSSC-0.2xSSC
5 buffer, 0.1% (w/v) SDS at a temperature of at least 55°C. Conditions for hybridisations and washes are well understood by one normally skilled in the art. For the purposes of further clarification only, reference to the parameters affecting hybridisation between nucleic acid molecules is found in Ausubel *et al.* (1992), which is herein incorporated by reference.

- 10 In an even more preferred embodiment of the invention, a hybridising nucleic acid molecule further comprises a sequence of nucleotides which is at least 40% identical to at least 10 contiguous nucleotides, preferably at least 50 contiguous nucleotides and more preferably at least 100 contiguous nucleotides, derived from any one of <400>1 or <400>3 or <400>5 or <400>7 or <400>8 or <400>9 or <400>11 or <400>13 or a complementary strand thereof, or
15 from a cDNA contained in any one or more of the deposited plasmids.

- In determining whether or not two nucleotide sequences fall within these percentage limits, those skilled in the art will be aware that it is necessary to conduct a side-by-side comparison or multiple alignment of sequences. In such comparisons or alignments, differences may arise
20 in the positioning of non-identical residues, depending upon the algorithm used to perform the alignment. In the present context, reference to a percentage identity between two or more nucleotide sequences shall be taken to refer to the number of identical residues between said sequences as determined using any standard algorithm known to those skilled in the art. For example, nucleotide sequences may be aligned and their identity calculated using the
25 BESTFIT programme or other appropriate programme of the Computer Genetics Group, Inc., University Research Park, Madison, Wisconsin, United States of America (Devereaux *et al.*, 1984).

- In an alternative embodiment, nucleotide sequences encoding EcR polypeptide subunits of
30 insect steroid receptors or fragments thereof and/or EcR partner proteins (USP polypeptides) of insect steroid receptor or fragments thereof, or USP polypeptides of insect juvenile hormone

- 25 -

receptor polypeptides, are amplified in the polymerase chain reaction. According to this embodiment, one or two or more nucleic acid "primer molecules" derived from a nucleotide sequence exemplified herein as <400>1 or <400>3 or <400>5 or <400>7 or <400>8 or <400>9 or <400>11 or <400>13 or a complementary strand thereof, or from a cDNA contained in any
5 one or more of the deposited plasmids, are annealed or hybridized to a nucleic acid "template molecule" which at least comprises a nucleotide sequence encoding a related genetic sequence or a functional part thereof, and nucleic acid molecule copies of the template molecule are amplified enzymatically using a thermostable DNA polymerase enzyme, such as *Taq* polymerase or *Pfu* polymerase, amongst others.

10

More particularly, one of the primer molecules comprises contiguous nucleotides derived from any one of <400>1 or <400>3 or <400>5 or <400>7 or <400>8 or <400>9 or <400>11 or <400>13, or alternatively, from a cDNA contained in any one or more of the deposited plasmids; and another of said primers comprises contiguous nucleotides complementary to
15 <400>1 or <400>3 or <400>5 or <400>7 or <400>8 or <400>9 or <400>11 or <400>13, or alternatively, from a cDNA contained in any one or more of the deposited plasmids, subject to the proviso that the first and second primers are not complementary to each other.

In a preferred embodiment, each nucleic acid primer molecule is at least 10 nucleotides in
20 length, more preferably at least 20 nucleotides in length, even more preferably at least 30 nucleotides in length, still more preferably at least 40 nucleotides in length and even still more preferably at least 50 nucleotides in length.

Furthermore, the nucleic acid primer molecules consists of a combination of any of the
25 nucleotides adenine, cytidine, guanine, thymidine, or inosine, or functional analogues or derivatives thereof which are at least capable of being incorporated into a polynucleotide molecule without having an inhibitory effect on the hybridisation of said primer to the template molecule in the environment in which it is used.

30 Furthermore, one or both of the nucleic acid primer molecules may be contained in an aqueous mixture of other nucleic acid primer molecules, for example a mixture of degenerate

- 26 -

primer sequences which vary from each other by one or more nucleotide substitutions or deletions. Alternatively, one or both of the nucleic acid primer molecules may be in a substantially pure form.

- 5 In a particularly preferred embodiment exemplified herein, two primer nucleotide sequences are used to amplify related sequences, said primers comprising the nucleotide sequences as set forth in any one of <400>15 to <400>20 inclusive. Even more preferably, the primers are used in the combination of (i) <400>15 and <400>16; or (ii) <400>17 and <400>18; or (iii) <400>19 and <400>20.

10

The nucleic acid template molecule may be in a recombinant form, in a virus particle, insect cell, bacteriophage particle, yeast cell, animal cell, or a plant cell. Preferably, the nucleic acid template molecule is derived from an insect species.

- 15 Those skilled in the art will be aware that there are many known variations of the basic polymerase chain reaction procedure. Such variations are discussed, for example, in McPherson *et al* (1991). The present invention extends to the use of all such variations in the isolation of variant insect steroid receptor-encoding genes or fragments thereof, and/or variant partner protein-encoding genes or fragments thereof to those exemplified herein.

20

- The isolated nucleic acid molecule of the present invention, including those sequences exemplified herein and any variants thereof, may be cloned into a plasmid or bacteriophage molecule, for example to facilitate the preparation of primer molecules or hybridisation probes or for the production of recombinant gene products. Methods for the production of such
- 25 recombinant plasmids, cosmids, bacteriophage molecules or other recombinant molecules are well-known to those of ordinary skill in the art and can be accomplished without undue experimentation. Accordingly, the invention further extends to any recombinant plasmid, bacteriophage, cosmid or other recombinant molecule comprising the nucleotide sequence set forth in any one of <400>1 or <400>3 or <400>5 or <400>7 or <400>8 or <400>9 or <400>11
- 30 or <400>13 or <400>15 to <400>20, or a complementary sequence, homologue, analogue or derivative thereof, or a cDNA contained in any one or more of the deposited plasmids.

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 27 -

The nucleic acid molecule of the present invention is also useful for developing genetic constructs which comprise and preferably, express, the EcR polypeptide subunit of the insect steroid receptor and/or the EcR partner protein (USP polypeptide) of the steroid receptor and/or the USP polypeptide of the juvenile hormone receptor, thereby providing for the
5 production of the recombinant polypeptides in isolated cells or transformed tissues.

Accordingly, a further aspect of the present invention provides a genetic construct comprising the subject isolated nucleic acid molecule encoding the insect steroid receptor polypeptide or a juvenile hormone receptor polypeptide, operably linked to a promoter sequence. Preferably,
10 the subject nucleic acid molecule is in an expressible format, such that it is possible to produce a recombinant polypeptide therefrom.

Reference herein to a "promoter" is to be taken in its broadest context and includes the transcriptional regulatory sequences of a classical genomic gene, including the TATA box
15 which is required for accurate transcription initiation in a eukaryotic cell, with or without a CCAAT box sequence or alternatively, the Pribnow box required for accurate expression in prokaryotic cells.

Promoters may be cell, tissue, organ or system specific, or may be non-specific. Using
20 specific promoters, the expression of a bioactive agent or other polypeptide encoded by a structural gene to which the promoter is operably connected may be targeted to a desired cellular site. For example, in transgenic animals such as sheep, it can be envisaged that cells of the transgenic animal may contain a gene encoding a steroid receptor, preferably a steroid receptor linked to an epidermal specific promoter and a separate gene encoding, for example,
25 epidermal growth factor (EGF) which is functionally linked to one or more insect hormone response elements and may or may not also be linked to epidermal specific promoter elements. On administration of the appropriate insect steroid hormone to the transgenic animal, the activated complex between the insect steroid receptor and insect steroid may bind to the one or more insect steroid hormone response element thereby inducing EGF production
30 solely in epidermal cells which may give rise to defleecing. It is to be understood that this aspect of the invention is independent of the degree of thermostability of the insect steroid

DocId:32858560

- 28 -

receptor The same principal applies to expression of any bioactive molecule or reporter molecule in a specific cell type which is regulated by a transactivating complex between a steroid receptor complex and an appropriate insect steroid.

- 5 In the present context, the term "promoter" is also used to describe a synthetic or fusion molecule, or derivative which confers, activates or enhances expression in a cell in response to an external stimulus. Accordingly, the promoter may include further regulatory elements (i.e. upstream activating sequences, enhancers and silencers) which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner. Preferred
10 promoters may contain copies of one or more specific regulatory elements, in particular steroid responsive elements (SREs) or hormone-responsive elements (HREs), to further enhance expression and/or to alter the spatial expression and/or temporal expression pattern.

- Reference herein to the term "steroid response element" shall be taken to refers to one or
15 more *cis*-acting nucleotide sequences present in a naturally-occurring or synthetic or recombinant gene the expression of which is regulated by an insect steroid, such as an ecdysteroid, for example ecdysone or ponasterone A, wherein said regulation of expression results from an direct or indirect interaction between a steroid receptor and said *cis*-acting nucleotide sequence response element. Exemplary insect steroid hormone response elements
20 include the ecdysone response element hsp27 (EcRE) and any other nucleotide sequence which is capable of binding ecdysteroid receptors or polypeptide subunits thereof or fragments or analogies thereof (such as associated with E75, E74 or other *Drosophila* early genes), as described for example by Riddihough and Pelham (1987).

- 25 For example, an SRE or a plurality of such elements may be operably linked to a promoter such as the polyhedron promoter, p10 promoter, MMTV promoter or SV40 promoter, to make transcription of a structural gene to which said promoter is operably connected responsive to the presence of a steroid bound to the insect receptor (which may act as a transcription factor). One or more insect SREs may be located within a promoter, and may replace sequences
30 within a selected promoter which confer responsiveness to hormones or other agents which regulate promoter activity. Where response elements are different they may lead to

- 29 -

preferential binding of different insect steroids or analogues thereof such that a promoter may be differentially regulated.

Particularly preferred SREs according to this embodiment include, but are not limited to, the
5 hsp27 ecdysone response element described by Riddihough and Pelham (1987) or the 13 base-pair palindromic core contained therein.

A promoter is usually, but not necessarily, positioned upstream or 5', of a structural gene, the expression of which it regulates. Furthermore, the regulatory elements comprising a promoter
10 are usually positioned within 2 kb of the start site of transcription of the gene.

Placing a gene or isolated nucleic acid molecule operably under the control of a promoter sequence means positioning said gene or isolated nucleic acid molecule such that its expression is controlled by the promoter sequence. Promoters are generally positioned 5'
15 (upstream) to the genes that they control. In the construction of heterologous promoter/structural gene combinations it is generally preferred to position the promoter at a distance from the gene transcription start site that is approximately the same as the distance between that promoter and the gene it controls in its natural setting, i.e., the gene from which the promoter is derived. As is known in the art, some variation in this distance can be
20 accommodated without loss of promoter function. Similarly, the preferred positioning of a regulatory sequence element with respect to a heterologous gene to be placed under its control is defined by the positioning of the element in its natural setting, i.e., the genes from which it is derived. Again, as is known in the art, some variation in this distance can also occur.

25

Those skilled in the art will recognise that the choice of promoter will depend upon the nature of the cell being transformed and when expression is required. Furthermore, it is well-known in the art that the promoter sequence used in the expression vector will also vary depending upon the level of expression required and whether expression is intended to be constitutive
30 or regulated.

667020 "0498600

- 30 -

For expression in eukaryotic cells, the genetic construct generally comprises, in addition to the nucleic acid molecule of the invention, a promoter and optionally other regulatory sequences designed to facilitate expression of said nucleic acid molecule. The promoter may be derived from a genomic clone which normally encodes the expressed protein or alternatively, it may
5 be a heterologous promoter derived from another genetic source. Promoter sequences suitable for expression of genes in eukaryotic cells are well-known in the art.

Suitable promoters for use in eukaryotic expression vectors include those capable of regulating expression in mammalian cells, insect cells such as Sf9 or Sf21. (*Spodoptera frugiperda*) cells,
10 yeast cells and plant cells. Preferred promoters for expression in eukaryotic cells include the p10 promoter, MMTV promoter, polyhedron promoter, the SV40 early promoter and the cytomegalovirus (CMV- IE) promoter, promoters derived from immunoglobulin-producing cells (see, United States Patent No 4,663,281), polyoma virus promoters, and the LTR from various retroviruses (such as murine leukemia virus, murine or Rous sarcoma virus and HIV), amongst
15 others (See, *Enhancers and Eukaryotic Gene Expression*, Cold Spring Harbor Press, New York, 1983, which is incorporated herein by reference). Examples of other expression control sequences are enhancers or promoters derived from viruses, such as SV40, Adenovirus, Bovine Papilloma Virus, and the like.

20 Wherein the expression vector is intended for the production of recombinant protein, the promoter is further selected such that it is capable of regulating expression in a cell which is capable of performing any post-translational modification to the polypeptide which may be required for the subject recombinant polypeptide to be functional, such as N-linked glycosylation. Cells suitable for such purposes may be readily determined by those skilled in
25 the art. By way of exemplification, Chinese hamster ovary (CHO) cells may be employed to carry out the N-terminal glycosylation and signal sequence cleavage of a recombinant polypeptide produced therein. Alternatively, a baculovirus expression vector such as the pFastBac vector supplied by GibcoBRL may be used to express recombinant polypeptides in Sf9 (*Spodoptera frugiperda*) cells, following standard protocols.

30

Numerous expression vectors suitable for the present purpose have been described and are

Accepted for publication

- 31 -

readily available. The expression vector may be based upon the pcDNA3 vector distributed by Medos Company Pty Ltd, Victoria, Australia, which comprises the CMV promoter and BGH terminator sequences for regulating expression of the recombinant polypeptide of the invention in a eukaryotic cell, when isolated nucleic acid sequences encoding same are inserted, in the
5 sense orientation relative to the CMV promoter, into the multiple cloning site of said vector. Alternatively, the SG5 expression vector of Greene *et al.* (1988), supplied by Stratagene, or the pQE series of vectors supplied by Qiagen are particularly useful for such purposes, as exemplified herein.

- 10 Examples of eukaryotic cells contemplated herein to be suitable for expression include mammalian, yeast, insect, plant cells or cell lines such as COS, VERO, HeLa, mouse C127, Chinese hamster ovary (CHO), WI-38, baby hamster kidney (BHK), MDCK, sf21 (insect) or Sf9 (insect) cell lines. Such cell lines are readily available to those skilled in the art.
- 15 The prerequisite for expression in prokaryotic cells such as *Escherichia coli* is the use of a strong promoter with an effective ribosome binding site. Typical promoters suitable for expression in bacterial cells such as *E. coli* include, but are not limited to, the *lacZ* promoter, temperature-sensitive λ_L or λ_R promoters, T7 promoter or the IPTG-inducible *tac* promoter. A number of other vector systems for expressing the nucleic acid molecule of the invention in
20 *E. coli* are well-known in the art and are described for example in Ausubel *et al* (1992).

Numerous vectors having suitable promoter sequences for expression in bacteria have been described, such as for example, pKC30 (λ_L ; Shimatake and Rosenberg, 1981), pKK173-3 (*tac*; Amann and Brosius, 1985), pET-3 (T7; Studier and Moffat, 1986) or the pQE series of
25 expression vectors (Qiagen, CA), amongst others.

Suitable prokaryotic cells include corynebacterium, salmonella, *Escherichia coli*, *Bacillus* sp. and *Pseudomonas* sp. amongst others. Bacterial strains which are suitable for the present purpose are well-known in the relevant art (Ausubel *et al*, 1992).

30

The genetic constructs described herein may further comprise genetic sequences

- 32 -

corresponding to a bacterial origin of replication and/or a selectable marker gene such as an antibiotic-resistance gene, suitable for the maintenance and replication of said genetic construct in a prokaryotic or eukaryotic cell, tissue or organism. Such sequences are well-known in the art.

5

Selectable marker genes include genes which when expressed are capable of conferring resistance on a cell to a compound which would, absent expression of said selectable marker gene, prevent or slow cell proliferation or result in cell death. Preferred selectable marker genes contemplated herein include, but are not limited to antibiotic-resistance genes such as
10 those conferring resistance to ampicillin, Claforan, gentamycin, G-418, hygromycin, rifampicin, kanamycin, neomycin, spectinomycin, tetracycline or a derivative or related compound thereof or any other compound which may be toxic to a cell.

The origin of replication or a selectable marker gene will be spatially-separated from those
15 genetic sequences which encode the recombinant receptor polypeptide or fusion polypeptide comprising same.

Preferably, the genetic constructs of the invention, including any expression vectors, are capable of introduction into, and expression in, an *in vitro* cell culture, or for introduction into,
20 with or without integration into the genome of a cultured cell, cell line and/or transgenic animal.

In a particularly preferred embodiment, the expression vector is selected from the group consisting of: pLcEcR (AGAL Accession No. NM99/04566); pLcUSP (AGAL Accession No. NM99/04565); pMpEcR (AGAL Accession No. NM99/04567); and pMpUSP (AGAL Accession
25 No. NM99/04568.).

A further aspect of the invention provides a cell comprising the subject isolated nucleic acid molecule which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide.

30

As used herein, the word "cell" shall be taken to refer to a single cell, or a cell lysate, or a

- 33 -

tissue, organ or whole organism comprising same, including a tissue, organ or whole organism comprising a clonal group of cells or a heterogenous mixture of cell types, which may be a prokaryotic or eukaryotic cell as described *supra*.

- 5 In a preferred embodiment, the cell of the present invention expresses the isolated or recombinant polypeptide encoded by the nucleic acid molecule.

In a preferred embodiment, the cell expresses a steroid receptor polypeptide or a fragment thereof which receptor is capable of binding to an insect steroid or analogue thereof or a
10 candidate insecticidally active agent to form an activated complex, and comprises a nucleic acid sequence encoding a bioactive molecule or a reporter molecule operably linked to one or more insect steroid response elements which on binding of the said activated complex promotes transcription of the nucleic acid sequence, wherein said cell on exposure to insect steroid or an analogue thereof, regulates expression of said bioactive molecule or allows
15 detection of said reporter molecule.

To produce the cells of the invention, host cells are transfected or co-transfected or transformed with nucleotide sequences containing the DNA segments of interest (for example, the insect steroid receptor gene, the recombinant steroid response elements, or both) by well-
20 known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas lipofection or calcium phosphate treatment are often used for other cellular hosts. See, generally, Sambrook *et al*, (1989); Ausubel *et al*, (1992); and Potrykus (1990). Other transformation techniques include electroporation, DEAE-dextran, microprojectile bombardment, lipofection, microinjection, and
25 others.

As used herein, the term "transformed cell" is meant to also include the progeny of a transformed cell.

- 30 In a further aspect of this invention, there is provided an animal (such as a mammal or insect), microorganism, plant or aquatic organism, containing one or more cells as mentioned above.

- 34 -

Reference to plants, microorganisms and aquatic organisms includes any such organisms.

In this embodiment of the invention, it is to be appreciated that administration of an insect steroid or an analogue thereof to an organism will induce expression of the desired bioactive molecule, such as a polypeptide, with attendant advantages. For example, an induced protein may have a therapeutic effect ameliorating a disease state or preventing susceptibility to disease or may modify in some way the phenotype of an organism to produce a desired effect. In humans, for example, cell transplants (such as liver cells) may under the action of insect steroids, produce desirable hormones such as insulin, growth hormone, growth factors and the like.

A further aspect of the invention provides a recombinant or isolated polypeptide comprising a steroid receptor polypeptide or juvenile hormone receptor polypeptide derived from an insect or a bioactive derivative or analogue thereof, wherein said polypeptide:

- (i) is selected from the list comprising EcR polypeptide of a steroid receptor, the partner protein (USP polypeptide) of a steroid receptor and the USP polypeptide of a juvenile hormone receptor; and
- (ii) comprises an amino acid sequence that is at least 40% identical to any one of the amino acid sequences set forth in <400>2, <400>4, <400>6, <400>10, <400>12 or <400>14;

wherein said polypeptide is substantially free of naturally-associated insect cell components.

In an alternative embodiment, the recombinant or isolated polypeptide comprising a steroid receptor polypeptide or juvenile hormone receptor polypeptide derived from an insect or a bioactive derivative or analogue thereof, wherein said polypeptide:

- (i) is selected from the list comprising EcR polypeptide of a steroid receptor, the partner protein (USP polypeptide) of a steroid receptor and the USP polypeptide of a juvenile hormone receptor; and
- (ii) comprises an amino acid sequence that is at least 40% identical to an amino acid sequence encoded by the cDNA present in any one of the plasmids deposited under AGAL Accession No. NM99/04565, NM99/04566, NM99/04567, or NM99/04568;

- 35 -

wherein said polypeptide is substantially free of naturally-associated insect cell components.

Reference herein to "substantially free of naturally associated insect cell components" refers to at least 80% purity, preferably more than 90% purity, and more preferably more than 95%
5 purity. Normally, purity is measured on a polyacrylamide gel with homogeneity determined by staining of protein bonds. Alternatively, high resolution may be necessary using HPLC or similar means. For most purposes, a simple chromatography column or polyacrylamide gel may be used to determine purity. A protein which is chemically synthesized or synthesized in a cell system different from an insect cell from which it naturally originates would be free of
10 naturally-associated insect cell components.

The present invention clearly provides for the isolation of EcR polypeptide subunits and EcR partner protein (USP polypeptide) subunits of ecdysteroid receptors and USP polypeptides of juvenile hormone receptors, from various organisms of the class *Insecta*, as described *supra*,
15 in addition to protozoa and helminth sources.

Insect steroid receptors are characterized by functional ligand-binding domains, and DNA-binding domains, both of which interact to effect a change in the regulatory state of a gene operably linked to the DNA-binding site of the holoreceptor or a polypeptide or polypeptide
20 fragment thereof. Thus, insect steroid receptors seem to be ligand-responsive transcription factors. Additionally, insect steroid receptors generally contain a DNA-binding domain (Domain C), and a ligand-binding domain (Domain E), separated and flanked by additional domains as identified by Krust *et al* (1986). The C domain preferably comprises a zinc-finger DNA-binding domain which is usually hydrophilic, having high cysteine, lysine and arginine
25 content. The E domain preferably comprises hydrophobic amino acid residues and is further characterized by regions E1, E2 and E3. The ligand-binding domain of the members of the insect steroid receptor superfamily is typically carboxyl-proximal, relative to a DNA-binding domain (Evans, 1988). The entire ligand-binding domain is typically between about 200 and 250 amino acids but is potentially shorter. This domain has the subregions of high homology,
30 designated the E1, E2 and E3 regions - which may be collectively referred to as the "E region".

Amino acid residues proximal to the C domain comprise a region initially defined as separate

DocId:34494566

- 36 -

A and B domains. Region D separates the more conserved domains C and E. Region D typically has a hydrophilic region whose predicted secondary structure is rich in turns and coils. The F region is carboxy proximal to the E region (see, Krust *et al, supra*).

- 5 The receptor polypeptides of the present invention exhibit at least a ligand-binding domain, as characterized by sequence homology to regions E1, E2 and E3. The ligand-binding domains of the present invention are typically characterized by having significant homology in sequence and structure to these three regions. Fragments of insect steroid receptors and partner proteins capable of binding insect steroids, and candidate insecticidally active compounds
- 10 comprise an E-region or a sufficient portion of the E-region to allow binding.

Preferably, the recombinant or isolated EcR polypeptide subunit of the insect steroid receptor or EcR partner protein (USP polypeptide) subunit of the steroid receptor or USP polypeptide of the juvenile hormone receptor as described herein is thermostable.

15

- By "thermostable" is meant that a stated integer does not exhibit reduced activity at bacterial, plant or animal physiological temperatures above about 28°C or above about 30°C. The thermostability of insect steroid hormone receptors also refers to the capacity of such receptors to bind to ligand-binding domains or regions and/or to transactivate genes linked to
- 20 insect steroid hormone response elements at bacterial, plant or animal physiological temperatures above about 28°C or above about 30°C.

- The present invention clearly extends to variants of said polypeptides, as described *supra*. The polypeptide may be substantially free of naturally associated insect cell components, or may
- 25 be in combination with a partner protein which associates with the insect steroid receptor so as to confer enhanced affinity for insect steroid response elements, enhanced affinity for insect steroids or analogues thereof. For Example, the amino acid sequences exemplified herein may be varied by the deletion, substitution or insertion of one or more amino acids.

- 30 In one embodiment, amino acids of a polypeptide exemplified herein may be replaced by other amino acids having similar properties, for example hydrophobicity, hydrophilicity, hydrophobic

Accepted for publication

P:\OPER\MRO\ECDYSONE.C\P - 1/7/99

- 37 -

moment, charge or antigenicity, and so on.

Substitutions encompass amino acid alterations in which an amino acid of the base polypeptide is replaced with a different naturally-occurring or a non-conventional amino acid residue. Such substitutions may be classified as "conservative", in which case an amino acid residue contained in the base polypeptide is replaced with another naturally-occurring amino acid of similar character, for example Gly↔Ala, Val↔Ile↔Leu, Asp↔Glu, Lys↔Arg, Asn↔Gln or Phe↔Trp↔Tyr.

10 Substitutions encompassed by the present invention may also be "non-conservative", in which an amino acid residue which is present in the base polypeptide is substituted with an amino acid having different properties, such as a naturally-occurring amino acid from a different group (eg. substituted a charged or hydrophobic amino acid with alanine), or alternatively, in which a naturally-occurring amino acid is substituted with a non-conventional amino acid.

Those skilled in the art will be aware that several means are available for producing variants of the exemplified EcR polypeptide subunit of the insect steroid receptor or EcR partner protein (USP polypeptide) subunit of the steroid receptor or USP polypeptide of the juvenile hormone receptor, when provided with the nucleotide sequence of the nucleic acid molecule which encodes said polypeptide, for example site-directed mutagenesis of DNA and polymerase chain reaction utilising mutagenised oligonucleotide primers, amongst others.

Such polypeptide variants which are capable of binding insect steroids clearly form part of the present invention. Assays to determine such binding may be carried out according to
25 procedures well known in the art.

One such variant polypeptide encompassed by the present invention comprises an "in-frame" fusion polypeptide between different regions of different insect receptor polypeptides. As exemplified herein, the present inventors have discovered that, by producing synthetic genes
30 in which various domains of a base insect steroid receptor-encoding nucleotide sequence derived from a first source are interchanged or substituted with similar sequences derived from

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 38 -

a second source (referred to as "domain swapping"), it is possible to modify the bioactivity of the insect steroid receptor encoded therefor. For example, the biological activity of the EcR polypeptide of the *L. cuprina* or *M. persicae* ecdysone receptor exemplified herein may be modulated by replacing portions of its C-terminal or N-terminal sequences with the equivalent domains from the EcR polypeptide of the *D. melanogaster* ecdysone receptor or alternatively, by swapping regions of the EcR polypeptides of the *L. cuprina* and *M. persicae* ecdysone receptors *per se*.

As a further refinement, such changes in biological function can similarly be effected by making specific changes (e.g. addition, substitution or deletion) to only those amino-acids within each domain that are critical for determining the relevant catalytic function (eg. ligand-binding activity, DNA binding site affinity, etc), such as by site-directed mutagenesis.

According to this embodiment, there is provided a synthetic EcR polypeptide subunit of a steroid receptor, and/or a synthetic EcR partner protein (USP polypeptide) subunit of a steroid receptor, and/or a synthetic USP polypeptide of a juvenile hormone receptor, or an analogue or derivative of said synthetic polypeptides, wherein said synthetic polypeptides comprise an amino acid sequence which has the following properties:

- (i) it differs in amino acid sequence or exhibits different biological properties to a naturally-occurring EcR polypeptide subunit of a steroid receptor, and/or a naturally-occurring EcR partner protein (USP polypeptide) subunit of a steroid receptor, and/or a naturally-occurring USP polypeptide of a juvenile hormone receptor;
- (ii) it comprises a first sequence of amino acids which are at least about 40% identical to a part of any one of <400>2, <400>4, <400>6, <400>10, <400>12 or <400>14, or at least about 40% identical to a part of an amino acid sequence encoded by any one of the deposited plasmids, linked covalently to a second sequence of amino acids derived from an EcR polypeptide subunit of a steroid receptor, EcR partner protein (USP polypeptide) subunit of a steroid receptor, or USP polypeptide of a juvenile hormone receptor, wherein said first and second sequences are derived from different genomic sources.

5

25

30 In an alternative embodiment, the recombinant receptor polypeptide of the invention is produced as an "in-frame" fusion polypeptide with a second polypeptide, for example a

P:\OPER\IMPROVEDYSONE.CIP - 1/7/99

- 40 -

detectable reporter polypeptide such as β -galactosidase, β -glucuronidase, luciferase or other enzyme, or a FLAG peptide, hapten peptide such as a poly-lysine or poly-histidine or other polypeptide molecule.

- 5 By "in-frame" means that a nucleotide sequence which encodes a first polypeptide is placed (i.e. cloned or ligated) in the same open reading frame adjacent to a nucleotide sequence which encodes a second polypeptide with no intervening stop codons there between, such that when the ligated nucleic acid molecule is expressed, a single fusion polypeptide is produced which comprises a sequence of amino acids corresponding to the summation of the individual
10 amino acid sequences of the first and second polypeptides.

In order to produce a fusion polypeptide, the nucleic acid molecule which encodes the polypeptide of the invention, or an analogue or derivative thereof, is cloned adjacent to a second nucleic acid molecule encoding the second polypeptide, optionally separated by a
15 spacer nucleic acid molecule which encodes one or more amino acids (eg: poly-lysine or poly histidine, amongst others), such that the first coding region and the second coding region are in the same open reading frame, with no intervening stop codons between the two coding regions. When translated, the polypeptide thus produced comprises a fusion between the polypeptide products of the first and second coding regions. Wherein a spacer nucleic acid
20 molecule is utilised in the genetic construct, it may be desirable for said spacer to at least encode an amino acid sequence which is cleavable to assist in separation of the fused polypeptide products of the first and second coding regions, for example a thrombin cleavage site.

- 25 A genetic construct which encodes a fusion polypeptide further comprises at least one start codon and one stop codon, capable of being recognised by the cell's translational machinery in which expression is intended.

Preferably, a genetic construct which encodes a fusion polypeptide may be further modified
30 to include a genetic sequence which encodes a targeting signal placed in-frame with the coding region of the nucleotide sequence encoding the fusion polypeptide, to target the

Accepted for publication

P:\OPERIMROLECDYSONE.CIP - 1/7/99

- 41 -

expressed recombinant polypeptide to the extracellular matrix or other cell compartment. More preferably, the genetic sequence encoding targeting signal is placed in-frame at the 5'-terminus or the 3'-terminus, but most preferably at the 5'-terminus, of the coding region of the nucleotide sequence which encodes the fusion polypeptide.

5

Methods for the production of a fusion polypeptide are well-known to those skilled in the art.

The recombinant EcR polypeptide subunit of the insect steroid receptor or EcR partner protein (USP polypeptide) subunit of the steroid receptor or USP polypeptide of the juvenile hormone
10 receptor may be purified by standard techniques, such as column chromatography (using various matrices which interact with the protein products, such as ion exchange matrices, hydrophobic matrices and the like), affinity chromatography utilizing antibodies specific for the protein or other ligands such as dyes or insect steroids which bind to the protein.

15 Wherein the recombinant polypeptide is expressed as a fusion polypeptide, it is also possible to purify the fusion polypeptide based upon its properties (eg size, solubility, charge etc). Alternatively, the fusion polypeptide may be purified based upon the properties of the non-receptor moiety of said fusion polypeptide, for example substrate affinity. Once purified, the fusion polypeptide may be cleaved to release the intact polypeptide of the invention.

20

Alternatively, proteins may be synthesized by standard protein synthetic techniques as are well known in the art.

In a preferred embodiment, the recombinant or isolated polypeptides of the invention are
25 provided as a precipitate or crystallized by standard techniques, preferably for X-ray crystal structure determination.

The three-dimensional structure of the polypeptide of the invention or a holoreceptor comprising same or a fragment of said polypeptide or holoreceptor is particularly useful for
30 identifying candidate insecticidal agents which mimic ligands that bind to said three-dimensional structure and/or modulate the ability of insect steroids to bind thereto and activate

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 42 -

the receptor (see, for example, Von Itzstein *et al.*, 1993; and Bugg *et al.*, 1993).

According to this embodiment, the EcR polypeptides of the invention or ligand binding domains thereof, or their complexes with EcR partner proteins or ligand binding domains thereof, which confer enhanced affinity for insect steroid response elements or partner proteins (USP polypeptides) or ligands, are particularly useful to model the three-dimensional structure of the receptor ligand-binding region. In this manner, insecticidal compounds may be produced which bind to, or otherwise interact with, the ligand-binding region of the receptor and/or preferably interfere with ligand binding. In the same way, compounds may be developed which have a potentiated interaction with the insect steroid receptor over and above that of the physiological insect steroid which binds to the receptor.

Accordingly, a still further aspect of the invention provides a method of identifying a candidate insecticidally-active agent comprising the steps of:

- 15 a) expressing a USP polypeptide of a juvenile hormone receptor or a fragment thereof which includes the ligand-binding region, optionally in association with an EcR polypeptide of a steroid receptor or ligand binding domain thereof, and optionally in association with an insect steroid or analogue thereof, so as to form a complex;
- b) purifying or precipitating the complex;
- 20 c) determining the three-dimensional structure of the ligand binding domain of the complex; and
- d) identifying compounds which bind to or associate with the three-dimensional structure of the ligand binding domain, wherein said compounds represent candidate insecticidally-active agents.

25

Standard procedures are used to determine the three dimensional structure of the receptor polypeptides of the invention, for example using X-ray crystallography and/or nuclear magnetic resonance analysis (see, for example, Bugg *et al.*, 1993; Von Itzstein *et al.*, 1993).

30 Insecticidally-active agents contemplated herein include synthetic chemicals that mimic one or more ligands of the holoreceptor or its polypeptide subunit, or the ligand-binding region of

- 43 -

said holoreceptor or subunit, thereby modulating binding of steroids to said holoreceptor or subunit. Preferred insecticidally-active agents include bisacylhydrazines, iridoid glycosides or other non-steroidal modulators of ecdysteroid receptors or insect juvenile hormone receptors. Additionally, because the EcR partner protein (USP polypeptide) subunits of insect steroid
5 receptors, and the USP polypeptides of insect juvenile hormone receptors, bind insect juvenile hormones, a sesquiterpenoid group of ligands that regulate developmental transitions in insects (see Jones and Sharp, 1997), compounds which interfere with the binding of juvenile hormone are also candidate insecticides.

10 A further aspect of the present invention provides a method of identifying a modulator of insect steroid receptor-mediated gene expression or insect juvenile hormone receptor-mediated gene expression comprising:

- (i) assaying the expression of a reporter gene in the presence of a recombinant or isolated insect steroid receptor polypeptide or a juvenile hormone receptor
15 polypeptide of the invention and a potential modulator; and
- (ii) assaying the expression of a reporter gene in the presence of a recombinant or isolated insect steroid receptor polypeptide or a juvenile hormone receptor polypeptide of the invention and without said potential modulator; and
- (ii) comparing expression of the reporter gene in the presence of the potential
20 modulator to the expression of a reporter gene in the absence of the potential modulator,

wherein said reporter gene is placed operably under the control of a steroid response element (SRE) to which said insect steroid receptor binds or a promoter sequence comprising said SRE.

25

In the present context, a "modulator" is a compound or molecule that agonises or antagonises the binding properties and/or biological activity of a receptor polypeptide or holoreceptor. Preferred modulators according to this embodiment include those synthetic compounds that are suitable for use as insecticidally-active agents described *supra*.

30

The reporter gene may be any gene, the expression of which may be monitored or assayed

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 44 -

readily. Preferably, the reporter gene is a structural gene that encodes a peptide, polypeptide or enzyme that is assayed readily by enzymic or immunological means, for example the β -galactosidase, β -glucuronidase, luciferase or chloramphenicol acetyltransferase (CAT) genes. Alternatively, the reporter gene may be a gene which encodes an immunologically-detectable
5 protein, for example a FLAG peptide, poly-lysine peptide or poly-histidine peptide.

Standard methods are used to assay the expression of the reporter gene.

This embodiment of the invention may be applied directly to the identification of potential
10 insecticidally-active compounds or alternatively, modified for such purposes by assaying for the binding (direct or indirect) of the recombinant or isolated insect steroid receptor polypeptide or a juvenile hormone receptor polypeptide of the invention to a steroid response element (SRE), rather than by assaying for reporter gene expression. According to this alternative embodiment, the binding assayed in the presence or absence of a potential
15 insecticidally-active compound is compared, wherein a difference in the level of binding indicates that the candidate compound possesses potential insecticidal activity.

In addition, substances may be screened for insecticidal activity by assessing their ability to bind, *in vivo* or *in vitro*, to the intact ecdysone receptor or alternatively, the ligand-binding
20 regions of the EcR polypeptide subunit of the ecdysone receptor (eg. <400>2 and/or <400>6 and/or <400>10) and/or the EcR partner protein (USP polypeptide) of the ecdysone receptor (eg. <400>4 and/or <400>12 and/or <400>14). Competition assays involving the native insect steroid may be employed to assess insecticidal activity.

25 The performance of this embodiment may, for example, involve binding the insect steroid receptor polypeptide to a support such as a plurality of polymeric pins, whereafter the polypeptide resident on the plurality of pins is brought into contact with candidate insecticidal molecules for screening. The molecules being screened may be isotopically labelled so as to permit ready detection of binding. Alternatively, reporter molecules may be utilized which bind
30 to the insect steroid receptor candidate molecule complex. Alternatively, compounds for screening may be bound to a solid support, such as a plurality of pins which are then reacted

P:\OPER\MRO\ECDSONE.CIP - 1/7/99

- 45 -

with the thermostable insect steroid receptor or complex with a partner protein. Binding may, for example, be determined again by isotopic-labelling of the receptor, or by antibody detection or use of another reporting agent.

- 5 In an alternative embodiment, insecticidally-active agent are identified using rational drug design, by expressing a USP polypeptide of a juvenile hormone receptor or a fragment thereof which includes the ligand-binding region, optionally in association with an EcR polypeptide of a steroid receptor or ligand binding domain thereof, and optionally in association with an insect steroid or analogue thereof, so as to form a complex, determining the three-dimensional
10 structure of the ligand binding domain of the complex, and identifying compounds which bind to or associate with the three-dimensional structure of the ligand binding domain, wherein said compounds represent candidate insecticidally-active agents.

- The methods described herein for identifying modulators of gene expression and insecticidal
15 compounds, may be performed using prokaryotic or eukaryotic cells, cell lysates or aqueous solutions.

- A further aspect of this invention accordingly relates to synthetic compounds derived from the three dimensional structure of EcR polypeptides and/or EcR partner protein (USP polypeptide)
20 subunits of insect steroid receptors, or fragments thereof, or insect steroid receptors or fragments thereof, or USP polypeptides of insect juvenile hormone receptors or fragments thereof, which compounds are capable of binding to said receptors which have the effects of either inactivating the receptors (and thus acting as antagonists) or potentiating the activity of the receptor.

25

By "derived from" it is meant that the compounds are based on the three dimensional structure of the aforementioned proteins, that is, synthesized to bind, associate or interfere with insect steroid binding or juvenile hormone binding.

- 30 The compounds may bind strongly or irreversibly to the ligand binding site or another region of the receptor or USP and act as agonists or antagonists of insect steroids, or juvenile

- 46 -

5

EXAMPLE 1

A 3110 base-pair Fsp1-HindIII fragment was excised from a cDNA encoding the EcR polypeptide subunit of the *D. melanogaster* ecdysone receptor (Koelle *et al.*, 1991), the excised
15 sequence comprising the complete 2634 base pair coding region and 214 base pairs of 5'-leader sequence and 258 base pairs of 3'- untranslated sequence. The fragment was ligated into the BamH1 site of the expression plasmid pSG5 (Greene *et al.*, 1988) to produce the expression plasmid pSV40-EcR, wherein expression of the EcR polypeptide subunit of the *Drosophila melanogaster* ecdysone receptor is placed operably under the control of the SV40
20 promoter sequence.

EXAMPLE 2

Construction of the reporter plasmid p(EcRE)₇-CAT

25 The reporter plasmid p(EcRE)₇-CAT was constructed by insertion of seven copies of the hsp27
ecdysone response element, containing a central 13 base pair palindromic ecdysone response
element (EcRE), derived from the hsp27 gene (Riddihough and Pelham, 1987) into the HindIII
site of the plasmid pMMTV-CAT (Hollenberg and Evans, 1988), 93 base pairs upstream of the
transcription start site of the MMTV promoter, thereby operably connecting expression of the
30 chloramphenicol acetyltransferase structural gene to regulation by an insect receptor which
binds to the hsp27 ecdysone response element.

- 47 -

EXAMPLE 3

Cell Culture and Transient Transfection

Chinese hamster ovary (CHO) cells were maintained in 50% (v/v) Dubbecco's modified Eagle's medium (DMEM) and 50% (v/v) Hamm F12 nutrient mixture (GIBCO) supplemented with 10% (v/v) foetal bovine serum. Transfection was carried out by the DNA-calcium phosphate co-precipitation method (Ausubel *et al*, 1992). One day before transfection with the plasmids described in Examples 1 and/or 2, or other expression plasmids, CHO cells were plated out at $5 - 8 \times 10^5$ cells per 6 cm diameter culture dish in the above DMEM/F12 medium. Three hours before the addition of the DNA-calcium phosphate co-precipitate, the cells were washed with phosphate buffered saline (PBS; Sambrook *et al*, 1989) and cultured in fresh DMEM plus 10% (v/v) foetal bovine serum. The cells were incubated in the presence of the co-precipitate for eighteen hours before excess DNA was removed by washing with PBS. The cells were then cultured for another day in DMEM/F12 supplemented with 10% (v/v) foetal bovine serum with or without added ponasterone A (PNA), before harvesting. Cells were washed with PBS, harvested by mechanical scraping in 0.25 M Tris-HCl (pH 7.8), and disrupted by three freeze-thaw cycles.

All transfections included, in addition to expression and reporter plasmids, a β -galactosidase-expressing plasmid designated pPgK-LacZ (McBurney *et al*, 1991), which served as an internal control for the efficiency of transfection, and pUC18 DNA in an amount sufficient to produce 10 μ g total DNA per culture dish.

The chloramphenicol acetyltransferase (CAT) and β -galactosidase activities encoded by the reporter genes present in the reporter plasmids were assayed as described in Sambrook *et al*, (1989). Cells that were co-transfected with p(EcRE) $_{\gamma}$ -CAT and pSV40-EcR clearly showed induction of CAT activity in the presence of PNA, showing 50 units of activity. Controls showed negligible activity.

We have observed that the ecdysone receptor can lead to stimulation of expression from an ecdysone responsive promoter in some cell types, for example in CHO cells, but not in CV-1

- 48 -

cells. Whilst not being bound by any theory or mode of action, this may reflect a cell-type specific distribution of at least one other transcription factor essential for ecdysone responsiveness. To determine cell types suitable for expressing reporter genes under the control of the steroid receptor of the present invention, the cell-type specificity of ecdysone-responsive gene expression is assayed in cell-free transcription lysates derived from several target cell lines. Additionally, by fractionating and/or isolating the nuclear proteins of cell lines that express the reporter genes and supplementing lysates derived from non-expressing cell lines with such nuclear protein fractions or isolated proteins, any essential auxiliary factors are defined and the genes encoding them cloned. Co-transfection of the receptor-encoding genes with genes encoding such auxiliary factors removes limitations imposed by cell-type restricted ecdysone responsiveness.

EXAMPLE 4

Testing the Effect of temperature on transient expression

To determine whether the *D. melanogaster* ecdysone receptor polypeptide is stable at physiological temperatures above about 30°C, CHO cells were transfected as described in Example 3, with the plasmid pSV40-EcR and the reporter plasmid p(EcRE)₇-CAT in the presence of PNA, at 30°C and 37°C.

Briefly, CHO cells were plated out at 37°C sixteen to twenty hours before transfection. After washing away the DNA, the cells were cultured for two hours in fresh medium with or without hormone and the dishes divided into duplicate sets. One set was cultured for another day at 37°C before harvesting for CAT and β-galactosidase assays. The other set was cultured for three days at 30°C before assaying enzyme activities. Results indicated a reduction in the fold-induction of gene expression regulated by the *D. melanogaster* ecdysone receptor polypeptide at 37°C, compared to the fold-induction at 30°C, as shown in Table 1.

- 49 -

EXAMPLE 5**Attempted screening of an *L. cuprina* genomic DNA library to isolate genes encoding the EcR polypeptide subunit of the *L. cuprina* ecdysone receptor**

- 5 A 627 bp Eco - Kpn I fragment encompassing the DNA-binding domain of the EcR polypeptide subunit of the *D. melanogaster* ecdysone receptor was isolated, radioactively labelled and used to screen a *L. cuprina* genomic library constructed in bacteriophage lambda (prepared by CSIRO, division of Entomology, Canberra, Australia). In the first round of screening, twenty-four regions of the plates showed potential positive hybridization to the *D. melanogaster* probe.
- 10 However, second-round screening of these 24 first round positive plaques failed to yield any plaque giving a reproducible positive signal when hybridized to the *D. melanogaster* probe.

TABLE 1

pSV40-EcR (μ g/dish)	PNA (μ M)	Fold-induction of expression	
		37°C	30°C
2.5	20	14X	35X
	100	59X	54X
0.5	20	8X	26X
	100	47X	33X
0.1	20	1.6X	25X
	100	9.0X	39X

EXAMPLE 6**Cloning and characterization of a cDNA molecule encoding the EcR polypeptide of the *L. cuprina* ecdysone receptor****25 Rationale for amplification primer design**

The nucleotide sequences of the primers Rdna3 (400>15) and Rdna4 (<400>16) were derived from the amino acid sequence conserved between the DNA-binding domains of the EcR polypeptide subunits of the *D. melanogaster* and *C. tentans* ecdysone receptors. However,

- 50 -

amino acid sequences homologous to two other members of the steroid receptor superfamily of *D. melanogaster*, *Drosophila* hormone receptor 3 (DHR3; Koelle, *et al.*, 1991) and *Drosophila* early gene (E75; Segraves and Hogness, 1990) were excluded from the primer designs, to reduce the possibility of amplifying the *L. cuprina* homologues of genes encoding
5 DHR3 and/or E75 by PCR.

Amplification primers and PCR conditions

A 105 base pair DNA fragment, encoding the DNA-binding domain of the EcR polypeptide subunit of the *L. cuprina* ecdysone receptor, was amplified from the *L. cuprina* genome by
10 PCR, by using the following degenerate primers:

Rdna3 (32mer with EcoRI site):

5'-CGGAATTCCGCCTCTGGTTA(C/T)CA(C/T)TA(C/T)AA(C/T)GC 3' (i.e. <400>15); and

Rdna4 (32mer with BamHI site):

15 5'-CGCGGATCC(G/A)CACTCCTGACACTTTCG(C/T)CTCA 3' (i.e. <400>16).

Amplification reactions employed *TaqI* DNA polymerase (Promega) and the following amplification conditions:

cycle 1: 97°C/5 minutes, 50°C hold; add polymerase 50°C/5 minutes;

20 cycles 2-3: 72°C/3 minutes, 94°C/1 minute, 50°C/1 minute;

cycles 4-43: 72°C/3 minutes, 94°C/1 minute, 55°C/1 minute;

cycle 44: 72°C/10 minutes.

To facilitate cloning of the amplified fragments for use as hybridisation probes, the 5' end of
25 primer Rdna3 contained an *EcoRI* site and the 5' end of primer Rdna4 contained a *BamHI* site. The amplified *L. cuprina* gene fragments were cloned into *pBluescript* SK+, following digestion using the enzymes *EcoRI* and *BamHI*, purification of the digested DNA by agarose gel electrophoresis and electro elution of the product band.

30 Hybridisation probe preparation

For probe preparation, the insert was cut out of the *pBluescript* SK+ vector using *EcoRI* and

- 51 -

BamHI, and ^{32}P -labelled using the GIGAprime DNA Labelling Kit (Bresatec Limited, Adelaide, Australia) essentially according to the manufacturer's instructions, except that random primers were replaced with the specific primers Rdna3 and Rdna4 (see above). Unincorporated label was removed by size exclusion chromatography over Biogel-P60 (Biorad Ltd, Sydney, Australia). The probe was used at 10^6 cpm/ml in hybridizations.

Construction and screening of *L. cuprina* cDNA libraries

Two independent *L. cuprina* cDNA libraries derived from late third instar *L. cuprina* larvae were prepared by random priming and oligo-dT priming respectively, and cloned into the *Eco*RI site of the *Lambda/ZapII* vector (Stratagene). The primary libraries generated were subsequently amplified according to the manufacturer's instructions, using standard protocols.

Both cDNA libraries generated are superior to existing *L. cuprina* libraries in terms of their phage titre (i.e. pfu/ml) and insert sizes (0.5 - 4 kbp in both cases). In particular, the primary oligo-dT primed library comprised 4.7×10^6 pfu, whilst the amplified oligo-dT primed library comprised 7.5×10^{10} pfu/ml; the primary random-primed library comprised 1.3×10^6 pfu, whilst the amplified random-primed library comprised 3.4×10^{10} pfu/ml.

The prepared cDNA libraries were screened by lifting 500,000 plaques from each library in duplicate on to Hybond N membranes (Amersham) and hybridizing same under low stringency conditions to the ^{32}P -labelled amplification product produced using the primers Rdna3 and Rdna4 (see above). In particular, hybridisations were performed for twenty four hours at 37°C in a hybridisation solution comprising 42% (w/v) formamide; 5 x SSPE solution; 5 x Denhardt's solution; and 0.1% (w/v) sodium dodecyl sulphate, as described essentially by Ausubel *et al*, (1992) and/or Sambrook *et al*. (1989). The membranes were then washed at 37°C in 2XSSC solution containing 0.1% (w/v) sodium dodecyl sulphate. Following washing, positive plaques were detected by autoradiography, using XOMAT-AR film (Kodak) for two to three days, at -70°C .

Two positive-hybridising plaques were obtained from screening of the random-primed library (containing cDNA inserts comprising 561 base pairs and 1600 base pairs in length,

Based upon the nucleotide sequence of the amplified DNA fragment, two authentic primers

- 53 -

were synthesized as follows:

Mdna1 (23mer): 5'- GCCTCGGGGTATCACTATAACGC -3' (i.e. <400>17); and

Mdna2 (23mer): 5'- GCACTCCTGACACTTTCGTCTCA -3' (i.e. <400>18).

5

Hybridisation probe preparation

For *M. persicae* probe preparation, the amplified 105 bp DNA insert was excised from the pBluescript SK+ vector using EcoRI and BamHI, and ³²P-labelled using the GIGAprime DNA Labelling Kit (BresaGen Limited, Adelaide, Australia) essentially according to the
10 manufacturer's instructions, except that random primers were replaced with the specific primers Mdna1 and Mdna2 (see above). Unincorporated label was removed by size exclusion chromatography over Biogel-P60 (Biorad Ltd, Sydney, Australia). The probe was used at 10⁶ cpm/ml in hybridizations.

15 Construction and screening of *M. persicae* cDNA libraries:

Two independent *M. persicae* cDNA libraries derived from late third instar *M. persicae* larvae were prepared by random priming and oligo-dT priming respectively, and cloned into the EcoRI site of the *Lambda/ZapII* vector (Stratagene). The primary libraries generated were subsequently amplified according to the manufacturer's instructions, using standard protocols.

20

Both cDNA libraries generated are superior to existing *M. persicae* libraries in terms of their phage titre (i.e. pfu/ml) and insert sizes (0.5 - 4 kbp in both cases). In particular, the primary oligo-dT-primed library comprised 1 x 10⁷ pfu, whilst the amplified oligo-dT primed library comprised 1 x 10¹⁰ pfu/ml; the primary random-primed library comprised 1 x 10⁶ pfu, whilst the
25 amplified random-primed library comprised 2 x 10¹¹ pfu/ml.

Additionally, a further cDNA library was produced in the Lambda ZAP Express insertion vector (Stratagene). To produce this library, cDNA derived from late third instar *M. persicae* larvae was prepared by oligo-dT priming and cloned directionally into EcoRI-XhoI digested vector
30 DNA. The primary library comprised 1 x 10⁶ pfu, whilst the amplified oligo-dT primed library comprised 1 x 10⁹ pfu/ml, with insert sizes in the range 0.5 - >4 kbp.

66 "P" 02434660

- 54 -

The random-primed *M. persicae* cDNA phage library was screened as described in Example 6, using the *M. persicae* hybridisation probe prepared as described above.

A single positive-hybridising plaque was isolated and sequenced according to standard 5 procedures. The nucleotide sequence of this clone is set forth herein as <400>5. This cDNA clone comprises a 585bp protein-encoding sequence which encodes the DNA-binding domain of a EcR polypeptide of a putative *M. persicae* ecdysone receptor. The amino acid sequence encoded by this partial cDNA clone is set forth herein as <400> 6.

10

EXAMPLE 8

Second attempt at cloning and characterization of a cDNA molecule encoding the EcR polypeptide of the *M. persicae* ecdysone receptor

15 Hybridisation probe preparation

Further hybridisation probes specific for the EcR polypeptide of the *M. persicae* ecdysone receptor were generated using PCR from the Lambda ZAPII oligo dT-primed library using primers AP1 and AP2. The forward primer AP1 was designed to anneal to nucleotide sequences of the partial cDNA (<400>5) encoding part of the first zinc finger motif present in 20 the DNA-binding domain. The reverse primer, AP2, was adapted from degenerate primers designed to anneal to nucleotide sequences complementary to those encoding an EcR ligand binding domain (Kamimura *et al.*, 1996). The nucleotide sequences of primers AP1 and AP2 are as follows:

- 25 Primer AP1: 5'- TCGTCCGGTTACCATTACAACGC -3' (<400>19); and
Primer AP2: 5'- TAGACCTTTGGC(A/G)AA(C/T)TC(A/G/C/T)ACAAT -3'(<400>20)

The PCR reaction mixture contained 4 µl of each primer (50 pm/µl), 5 µl of deoxynucleotide triphosphate mix (2mM), 1 µl of aphid oligo dT primed Lambda ZAPII cDNA library, 1 µl of 30 recombinant *Pfu* DNA Polymerase (5 units/µl, Stratagene®), 5 µl of 10x *Pfu* buffer (Stratagene®) and 30 µl of MilliQ water. The *Pfu* polymerase was used in this reaction

2020 "043166"

- 55 -

because it possesses proof-reading activity, which reduces the possibility of misincorporation of nucleotides. The PCR conditions included 42 cycles, each cycle comprising annealing at 55°C, extension at 72°C and melting at 94°C.

- 5 The major amplification product obtained in this reaction was gel-purified, kinased and ligated into the *Sma*I site of pUC18.

To screen *M. persicae* cDNA libraries, the cloned amplification product was digested to generate two non-overlapping probes, designated "EcR probe 1" (i.e. <400>7) and "EcR
10 probe 2" (i.e. <400>8). In this regard, digestion of the cloned product with *Sph*I produced a DNA fragment comprising a nucleotide sequence specific for a region encoding the DNA-binding domain (EcR probe 1; <400> 7), whilst digestion with *Sph*I/*Eco*RI produced a DNA fragment comprising a nucleotide sequence having homology to a region encoding a putative linker domain, designated domain D, and the 5'-end of a putative hormone-binding domain,
15 present in the EcR polypeptide of the insect ecdysone receptors (EcR probe 2, <400> 8).

EcR probe 1 and EcR probe 2 were labelled with [α -³²P]dATP in a reaction catalysed by Klenow fragment. All reagents were components of a GIGAprime DNA labelling kit (BresaGen Limited, Adelaide, Australia), except that the random primers were replaced with specific
20 oligonucleotides synthesized to be complementary to the ends of EcR probe 1 and EcR probe 2.

Screening of *M. persicae* cDNA libraries

480,000 plaques from the oligo dT primed Lambda Zap Express cDNA library (Example 7)
25 were screened as described above, using EcR probe 1. This approach yielded about 300 positive clones. Positive-hybridising clones were pooled and rescreened separately using EcR probe 1 and EcR probe 2, on duplicate lifts. Only four plaques were identified which hybridised to both probes. One of these was found by sequencing to contain a full-length cDNA encoding the EcR polypeptide of the *M. persicae* ecdysone receptor. The nucleotide sequence of the
30 open reading frame of this cDNA is set forth herein as <400> 9. The derived amino acid sequence of the EcR polypeptide subunit of the *M. persicae* ecdysone receptor encoded by

- 56 -

this open reading frame is set out in <400> 10.

EXAMPLE 9

In vivo function of recombinant EcR polypeptides of the

5 *L. cuprina* ecdysone receptor

Construction of plasmid pF3

Plasmid pF3 was constructed in four steps as follows:

First, plasmid p5S1, comprising the full-length cDNA encoding the EcR polypeptide of the *L.*
10 *cuprina* ecdysone receptor, was digested with *EarI* and a 3' *EarI* cDNA fragment thus
generated, encoding the C-terminal end of the EcR polypeptide of the *L. cuprina* ecdysone
receptor, was end-filled and sub-cloned into the *HindIII* site of pUC19, to construct plasmid
pEAR. In plasmid pEAR, the 3' end of the cDNA was oriented towards the *KpnI* site of the
pUC19 vector.

15 Second, plasmid p5S1 was also digested separately with:

- (1) *ApoI* and *PstI*, to isolate the 5' end of the cDNA as a 179 bp fragment (fragment A);
- (2) *PstI* and *SpeI*, to isolate a 1650 bp cDNA fragment (fragment B); and
- (3) *SpeI* and *BglII*, to isolate a 203 bp fragment (fragment C).

20 Third, plasmid pEAR was digested with *BglII* and *KpnI*, to isolate the 3' end of the cloned cDNA
fragment therein as a 313 bp fragment (fragment D).

Fourth, DNA fragments A, B, C and D were each isolated by agarose electrophoresis and
25 ligated together into pBluescriptSK+, which had been digested with *EcoRI* and *KpnI*, to
produce plasmid pF3.

Plasmid pF3 thus contains the complete open reading frame of the cDNA encoding the EcR
polypeptide of the *L. cuprina* ecdysone receptor, as a 2368 bp fragment located between two
30 *BamHI* sites.

667020 "02434660

- 57 -

Construction of plasmid pSGLcEcR and plasmid pLcK8

Plasmid pSGLcEcR was constructed by cloning the 2368 bp *Bam*HI fragment from pF3, into the *Bam*HI site of the mammalian expression vector pSG5 (Stratagene). Plasmid pLcK8 is a clone of pSGLcEcR.

5

Construction of plasmid pSGDmEcR

Plasmid pSGDmEcR is identical to plasmid pSV40-EcR (Example 1) comprising the EcR polypeptide of the *D. melanogaster* ecdysone receptor placed operably under control of the SV40 promoter.

10

Transfection of CHO cells

CHO cells were co-transfected with a mixture comprising the following DNAs, lysed and assayed for CAT and β -galactosidase enzyme activity, as described in the preceding Examples:

- 15 (1) one of the expression plasmids designated pSGDmEcR, or pSGLcEcR, or the parental expression plasmid pSG5 as a negative control, at a concentration of 1 μ g/ml; and
- (2) the CAT reporter plasmid p(EcRE)₃CAT at a concentration of 1 μ g/ml; and
- (3) an independent LacZ reporter plasmid, pPGKLacZ, at a concentration of 1 μ g/ml,
- 20 included as a control to monitor transfection efficiency .

CAT reporter gene expression was induced with 10 μ M or 50 μ M Muristerone A. In control samples, cells received only the carrier ethanol in place of Muristerone A.

ELISA was used to quantify the synthesis of CAT and β -galactosidase enzymes, in extracts

25 of cells forty eight hours after transfection. Account was taken of the variation between experiments, by normalizing the level of CAT enzyme to the level of β -galactosidase enzyme present in the same extract. Fold induction represents the normalized values for CAT gene expression in cells transfected with pSGDmEcR, pSGLcEcR or pSG5 in the presence of hormone divided by the normalized values for CAT gene expression in cells transfected with

30 the same plasmid but in the absence of hormone. The average values of three independent experiments are shown in Figure 1 and the error bars indicate standard error of the mean.

- 58 -

Data shown in figure 1 indicate that the EcR polypeptide of the *L. cuprina* ecdysone receptor from Example 3 is biologically active *in vivo*. CAT induction is observed at both 50 μ M and 10 μ M steroid (Muriesterone A), with about 30 and 15 fold induction respectively. In view of the *in vivo* activity of the EcR polypeptide of the *L. cuprina* ecdysone receptor obtained according to this protocol, potential insecticidal substances acting by interaction with an insect steroid receptor, such as an ecdysone receptor, are screened by addition of the substances to the *in vivo* assay described herein. Substances are added in an amount from 0.05 μ M to 100 μ M. Candidate insecticidal compounds are identified by their ability to modulate the reporter gene expression which results from trans-activation by the EcR polypeptide of the *L. cuprina* ecdysone receptor.

EXAMPLE 10

Chimeric EcR polypeptides of insect ecdysone receptors

Chimeric ecdysone receptors comprising regions derived from EcR polypeptides of ecdysone receptors of different species are produced and assayed for enhanced activity. In a particularly preferred embodiment, a chimeric ecdysone receptor is produced using the EcR polypeptides of the *D. melanogaster*, *M. persicae* and *L. cuprina* ecdysone receptors.

In one exemplification of this embodiment, plasmids pSGLD and pSGDL are produced comprising coding regions derived from the EcR polypeptides of the *D. melanogaster* and *L. cuprina* ecdysone receptors. In plasmid pSGLD, the 5'-end of the open reading frame of the *D. melanogaster* sequence, encoding the N-terminal portion of the EcR polypeptide of the *D. melanogaster* ecdysone receptor to the end of the DNA-binding domain of said polypeptide, is fused to the 3'-end of the open reading frame of the *L. cuprina* sequence, encoding the C-terminal portion of the EcR polypeptide of the *L. cuprina* ecdysone receptor, from the D domain and hormone-binding domain to the carboxyl terminus. In plasmid pSGDL, the 5'-end of the open reading frame of the *L. cuprina* sequence, encoding the N-terminal portion of the EcR polypeptide of the *L. cuprina* ecdysone receptor to the end of the DNA-binding domain of said polypeptide, is fused to the 3'-end of the open reading frame of the *D. melanogaster* sequence, encoding the C-terminal portion of the EcR polypeptide of the *D. melanogaster* ecdysone receptor, from the D domain and hormone-binding domain to the carboxyl terminus.

- 59 -

As shown in Figure 2, chimeric EcR polypeptides of *L. cuprina* and *D. melanogaster* ecdysone
5 receptors, comprising fusion polypeptides between the DNA-binding domains and hormone-
binding domains of the base *L. cuprina* and *D. melanogaster* polypeptides, exhibit bioactivity
when measured in the CAT assay described above. Significant bioactivity of the chimeric EcR
polypeptides encoded by plasmids pSGLD and pSGDL, comparable to the bioactivity of the
D. melanogaster base EcR polypeptide, is observed at both 10 μ M and 50 μ M concentrations
10 of Muristerone A.

Isolation and characterisation of a full-length cDNA encoding the EcR partner protein (USP polypeptide) of the *L. cuprina* ecdysone receptor

15 The EcR partner protein (USP polypeptide) subunit of the *L. cuprina* ecdysone receptor also functions alone as a USP polypeptide of the *L. cuprina* juvenile hormone receptor. A cDNA encoding both receptor polypeptide activities was isolated using PCR and hybridisation as follows.

20 Hybridisation probe preparation

A 150 base-pair probe, specific for genetic sequences encoding the EcR partner protein (USP polypeptide) subunit of insect ecdysone receptors and/or the USP polypeptide subunit of insect juvenile hormone receptors (<400>13), was isolated by PCR from *L. cuprina* genomic DNA using the degenerate primers described by Tzertzinis *et al.* (1994). The PCR reaction conditions were as described in Example 6, except that *Pfu* polymerase was used in place of *TaqI* polymerase.

The amplified DNA fragment was sub-cloned into *Eco*RI and *Cla*I double-digested *p*Bluescript SK+ vector (Stratagene), after double-digestion of the fragment using the enzymes *Eco*RI and *Cla*I, purification of the amplified fragment by agarose gel electrophoresis, and electro elution of the product band. The nucleotide sequence of the probe was obtained using the USB

- 60 -

Sequenase version 2.0 Kit (<400> 13).

For probe preparation, the amplified *L. cuprina* DNA fragment was excised from the vector using *EcoRI* and *SaI*, gel purified and ³²P-labelled using the GIGAPrime DNA Labelling Kit 5 (BresaGen Limited, Adelaide, Australia) essentially according to the manufacturer's instructions except that random primers were replaced with the two degenerate primers described by Tzertzinis *et al.* (1994) (see above). Unincorporated label was removed by size exclusion chromatography over Biogel-P60 (Biorad Ltd, Sydney, Australia). The probe was used at 10⁶ cpm/ml in hybridizations.

10

Screening of *L. cuprina* cDNA libraries

The *L. cuprina* cDNA library described above (Example 6) was screened with the amplified probe as described in Example 6. The nucleotide sequence of the full-length open reading frame of this cDNA molecule and amino acid sequence therefor, are set forth herein as <400>

15 3 and <400> 4, respectively.

EXAMPLE 12

Isolation and characterisation of a partial cDNA encoding the EcR partner protein (USP polypeptide) of the *M. persicae* ecdysone receptor

20

The EcR partner protein (USP polypeptide) subunit of the *M. persicae* ecdysone receptor also functions alone as a USP polypeptide of the *M. persicae* juvenile hormone receptor. To isolate a partial cDNA encoding both receptor polypeptide activities, a 140 bp probe was amplified 25 from *M. persicae* genomic DNA, by PCR, using the two degenerate primers described by Tzertzinis *et al.* (1994) (see preceding Example). The PCR reaction conditions were as described in Example 6, except that *Pfu* polymerase was used in place of *TaqI* polymerase.

The amplified DNA fragment was sub-cloned into *EcoRI* and *ClaI* double-digested *pBluescript* 30 SK+ vector (Stratagene), after double-digestion of the fragment using the enzymes *EcoRI* and *ClaI*, purification of the amplified fragment by agarose gel electrophoresis, and electro elution

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 61 -

of the product band.

The nucleotide sequence of the insert in the pBluescript SK+ vector was obtained using automated fluorescent dye terminator sequencing (SUPAMAC, Sydney Australia).

5

Hybridisation probe preparation and library screening

For probe preparation the amplified *M. persicae* DNA insert was cut out of the pBluescript+ vector with *EcoRI* and *SaII*, gel purified and ³²P-labelled using the GIGAprime DNA Labelling Kit (Bresatec Limited, Adelaide, Australia) essentially according to the manufacturer's instructions except that random primers were replaced with the degenerate primers described by Tzertzinis *et al.* (1994) (see preceding Example). Unincorporated label was removed by size exclusion chromatography over Biogel-P60 (Biorad Ltd, Sydney, Australia). The probe was used at 10⁶ cpm/ml in hybridizations to screen the *M. persicae* cDNA library as described in Examples 7 and 8.

15

The positive-hybridising clones were plaque-purified and sequenced using standard procedures as described herein. The nucleotide sequence of the open reading frame of the full-length cDNA encoding the partner protein (USP polypeptide) subunit of the *M. persicae* ecdysone receptor or the USP polypeptide of the *M. persicae* juvenile hormone receptor is set forth herein as <400> 11. The derived amino acid sequence of this open reading frame is set forth as <400>12.

EXAMPLE 13

A construct for the baculovirus-directed co-expression of functional ligand-binding regions of the EcR polypeptide and partner protein (USP polypeptide) of the *D. melanogaster* ecdysone receptor

A vector was prepared to facilitate the baculovirus-directed co-expression of ligand-binding regions derived from the EcR polypeptide and partner protein (USP polypeptide) of the *D. melanogaster* ecdysone receptor, the protein products of which associate on co-expression to form a functional hormone-binding complex. The associated proteins are then used in high

667020 "04794660

- 62 -

through-put assays or three-dimensional structural analysis. We have found that the ligand-binding domain, together with most of the linker domain of the EcR polypeptide subunit and of the EcR partner protein (USP polypeptide), are sufficient to associate to form a functional hormone-binding complex.

5

1. Isolation of the ligand-binding region and linker region of the EcR polypeptide of the *D. melanogaster* ecdysone receptor.

A Sac I- HindIII fragment encoding most of the linker (domain D) and all of the ligand-binding domain (domains E and F) of the EcR polypeptide of the *Drosophila melanogaster* ecdysone
10 receptor was excised from a plasmid comprising DNA encoding the complete EcR polypeptide (Koelle *et al.* 1991). The excised fragment was cloned into SacI - HindIII-digested expression vector pQE31(Qiagen), to produce the plasmid vector pQE31DmECR.

2. Construction of a baculovirus expressing the linker regions of EcR and USP polypeptides

15 A baculovirus was constructed for the co-expression in insect cells of:

- (i) a cDNA region comprising a nucleotide sequence which encodes at least the ligand-binding domain and much of the linker domain of the EcR polypeptide of the *D. melanogaster* ecdysone receptor isolated as described at paragraph (1) above; and
- (ii) a cDNA region comprising a nucleotide sequence which encodes at least the
20 ligand-binding domain and much of the linker domain of the partner protein (USP polypeptide) of the *D. melanogaster* ecdysone receptor.

To produce this baculovirus, a EcoR I - HindIII fragment was excised from pQE31DmECR, said fragment encoding an oligo-His tag, and most of the linker domain, together with all of the
25 ligand-binding domain of EcR polypeptide. This EcoR I - HindIII fragment was ligated into EcoR I - HindIII cleaved pFastBacDUAL, to produce the plasmid pDmEcR.DUAL. To insert gene sequences specific for the partner protein (USP polypeptide), the HindII - NsiI fragment encoding most of the linker and all of the ligand-binding domain of the partner protein (USP polypeptide) was excised from a full-length cDNA clone in plasmid pZ7-1 (supplied by Vince
30 Henrich) and ligated into NcoI - NsiI cleaved pDmEcR.DUAL. A nucleotide sequence encoding a "FLAG" peptide was subsequently incorporated upstream of, and in the same reading frame

 09440349460
667020"0249460

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 63 -

as, the nucleotide sequence encoding the linker and ligand-binding regions of the partner protein (USP polypeptide), by ligation into the unique *Sma*I site, thereby producing the plasmid pDmEcR.USP.DUAL. Plasmids containing the FLAG-encoding nucleotide sequence in the correct orientation were selected by nucleotide sequence determination.

5

The segment of pDmEcR.USP.DUAL which encodes the tagged linker and ligand-binding regions of the EcR polypeptide and partner protein (USP polypeptide) sequences, placed operably under the control of polyhedrin and p10 promoters, respectively, was recombined into a baculovirus genome, by employing the Tn7 transposition system (Luckow *et al*, (1993). The

10 polypeptide products were then co-expressed in insect Sf21 and Sf9 cells, where they associated into a functional complex.

Expression of the tagged linker and ligand-binding regions of the EcR polypeptide and partner protein (USP polypeptide) sequences was examined by immunoblot analysis of extracts

15 derived from insect Sf21 cells infected with the recombinant baculovirus, employing antibodies directed against the oligo-His and FLAG tags. This analysis detected bands on immunoblot analysis of approximately the predicted sizes for the expressed tagged linker and ligand-binding regions of the EcR polypeptide and partner protein (USP polypeptide).

20 The protein detected by anti-oligo-His-antibodies was enriched by affinity purification on nickel-NTA resin (Qiagen), and the FLAG-labelled protein was affinity-purified using FLAG M2 Affinity Gel (Kodak). It was further demonstrated that the oligo-His-tagged EcR polypeptide and the FLAG-tagged EcR partner protein (USP polypeptide) bound as a hetero-oligomeric complex to FLAG M2 Affinity Gel (Kodak).

25

Furthermore, binding assays, performed using a modification of the method of Yund *et al* (1978), demonstrated a highly-significant increase in the binding of the a labelled ecdysone analogue, [³H]ponasterone A, in cells infected by the recombinant baculovirus, compared to the binding observed for the naturally-occurring ecdysone holoreceptor in *L. cuprina* embryos.

30 In contrast, cells infected by a control virus displayed neither antibody-positive bands on western analysis, nor specific binding of [³H] ponasterone A, above background levels. These

- 64 -

data indicate correct folding and association of the variant polypeptides comprising the linker and ligand-binding regions of the *D. melanogaster* EcR polypeptide and *D. melanogaster* partner protein (USP polypeptide). The correctly-folded and associated complex formed by the truncated EcR polypeptide and truncated EcR partner protein (USP polypeptide), is used for

5 X-ray and NMR structural analysis and for high-throughput screens.

EXAMPLE 14

Construct for the baculovirus-directed co-expression of functional ligand-binding regions of the EcR polypeptide and partner protein (USP polypeptide) of the *L.*

10 *cuprina* ecdysone receptor

A vector for the baculovirus-directed co-expression of ligand-binding domains derived from the EcR polypeptide and partner protein (USP polypeptide) of the *L. cuprina* ecdysone receptor was prepared essentially as described in the preceding Example.

15

1. Isolation of the ligand-binding region and linker region of the EcR polypeptide of the *L. cuprina* ecdysone receptor.

A *SphI* – *KpnI* fragment encoding most of the linker (domain D) and all of the ligand-binding domain (domains E and F) of the EcR polypeptide of the *L. cuprina* ecdysone receptor was

20 excised from a cDNA clone encoding the complete EcR polypeptide and cloned into the *SphI* – *KpnI* cleaved expression vector pQE32 (Qiagen), to produce the plasmid pQE32LcEcR.

2. Isolation of the ligand-binding region and linker region of the partner protein (USP polypeptide) of the *L. cuprina* ecdysone receptor.

25 A DNA fragment encoding most of the linker domain and all of the ligand-binding domain of the partner protein (USP polypeptide) of the *L. cuprina* ecdysone receptor was sub-cloned to produce the plasmid pBLU1.

3. Construction of a baculovirus expressing the linker regions of *L. cuprina* EcR and USP

30 polypeptides

A baculovirus was constructed for the co-expression in insect cells of:

- 65 -

- (i) a cDNA region comprising a nucleotide sequence which encodes at least the ligand-binding domain and much of the linker domain of the EcR polypeptide of the *L. cuprina* ecdysone receptor isolated as described at paragraph (1) above; and
- (ii) a cDNA region comprising a nucleotide sequence which encodes at least the
5 ligand-binding domain and much of the linker domain of the partner protein (USP polypeptide) of the *L. cuprina* ecdysone receptor isolated as described at paragraph (2) above.

To produce this baculovirus, a *EcoR* I – *Pst*I fragment derived from plasmid pQE32LcEcR,
10 encoding an oligo-His tag and most of the linker domain together with all of the ligand-binding domain of the *L. cuprina* EcR polypeptide was ligated into *EcoR*I– *Pst*I cleaved pFastBac.DUAL, to produce the plasmid pLcEcR.DUAL. An *Avall*–*EcoR*V fragment, encoding most of the linker and all of the ligand-binding domain of *L. cuprina* partner protein (USP polypeptide) was excised from plasmid pBLU1 and ligated, together with a "FLAG" encoding
15 sequence into the *Pvu*II site of pLcEcR.DUAL, to produce plasmid pLcEcR.USP.DUAL.

The segment of pLcEcR.USP.DUAL which encodes the tagged linker and ligand-binding regions of the EcR polypeptide and partner protein (USP polypeptide) sequences, placed operably under the control of polyhedrin and p10 promoters, respectively, was recombined into
20 a baculovirus genome, by employing the Tn7 transposition system (Luckow *et al*, (1993). The polypeptide products were then co-expressed in insect Sf21 and Sf9 cells, where they associated into a functional complex.

Expression was examined by immunoblot analysis. Antibodies directed against oligo-His and
25 FLAG tags detected bands on immunoblot analysis of approximately the predicted sizes for the expressed EcR and USP polypeptide regions respectively, in extracts from insect Sf21 cells infected with the recombinant baculovirus. The protein detected by anti-oligo-His was greatly enriched utilising a nickel-NTA resin (Qiagen) and the FLAG-labelled protein purified on FLAG M2 Affinity Gel (Kodak). It was also demonstrated by immunoblot analysis that oligo-
30 His-tagged *L. cuprina* truncated EcR polypeptides and FLAG-tagged *L. cuprina* truncated EcR partner protein (USP polypeptide) bind as a hetero-oligomeric complex to FLAG M2 Affinity

P:\OPER\MRO\ECDYSONE\CIP - 1/7/99

- 66 -

Gel (Kodak).

Furthermore, binding assays, carried out by a modification of the method of Yund *et al* (1978), demonstrated a highly-significant increase in the binding of the tritiated ecdysone analogue, ponasterone A, in cells infected by recombinant virus indicating correct folding and association of the two protein subunits (Figure 3), greater than that of the ecdysone holoreceptor in *L. cuprina* embryos. Cells infected by a control virus displayed neither antibody-positive bands on western analysis nor specific binding of tritiated hormone above background.

- 10 Expression of the tagged linker and ligand-binding regions of the *L. cuprina* EcR polypeptide and partner protein (USP polypeptide) sequences was examined by immunoblot analysis of extracts derived from insect Sf21 cells infected with the recombinant baculovirus, employing antibodies directed against the oligo-His and FLAG tags. This analysis detected bands on immunoblot analysis of approximately the predicted sizes for the expressed tagged linker and
- 15 ligand-binding regions of the *L. cuprina* EcR polypeptide and partner protein (USP polypeptide).

The protein detected by anti-oligo-His-antibodies was enriched by affinity purification on nickel-NTA resin (Qiagen), and the FLAG-labelled protein was affinity-purified using FLAG M2 Affinity

20 Gel (Kodak).

Furthermore, binding assays, performed using a modification of the method of Yund *et al* (1978), demonstrated a significant increase in the binding of the labelled ecdysone analogue, [³H] ponasterone A, in cells infected by the recombinant baculovirus, compared to the binding

25 observed for the naturally-occurring ecdysone holoreceptor in *L. cuprina* embryos (Figure 3). In contrast, cells infected by a control virus displayed neither antibody-positive bands on western analysis, nor specific binding of [³H] ponasterone A, above background levels.

These data indicate correct folding and association of the variant polypeptides comprising the linker and ligand-binding regions of the *L. cuprina* EcR polypeptide and *L. cuprina* partner

30 protein (USP polypeptide). The correctly-folded and associated complex formed by the

- 67 -

truncated EcR polypeptide and truncated EcR partner protein (USP polypeptide), is used for X-ray and NMR structural analysis and for high-throughput screens.

EXAMPLE 15

5 **A construct for the expression of the ligand-binding region of the USP polypeptide of the *L. cuprina* juvenile hormone receptor**

The donor plasmid pLcEcR.USP.DUAL (Example 14) was digested with *Bss*HI and *Pst*I to remove the *L. cuprina* EcR polypeptide-encoding segment therein, thereby leaving the tagged
10 linker and ligand-binding regions of the *L. cuprina* USP polypeptide-encoding nucleotide sequence. The digested plasmid was blunt-ended using T4 DNA polymerase and Klenow polymerase, isolated by gel purification, and finally re-ligated to produce the plasmid pLc.USP.SINGLE.

15 To produce recombinant baculovirus capable of expressing the tagged linker and ligand-binding regions of the USP polypeptide, the segment of pLc.USP.SINGLE encoding this polypeptide and the p10 promoter sequence to which said segment is operably connected, is recombined into a baculovirus genome employing the Tn7 transposition system (Luckow *et al.*, 1993). The polypeptide product is then expressed to form a functional juvenile hormone-
20 binding polypeptide and preferably, a modulator of a juvenile hormone receptor. The correctly-folded truncated USP polypeptide is used for X-ray and NMR structural analysis and for high-throughput screens.

EXAMPLE 16

25 ***In-vitro* Screening for the Detection of Insecticidal Compounds**

The EcR partner protein (USP polypeptide) of the insect ecdysone receptor and USP polypeptide of the insect juvenile hormone receptor of the present invention, optionally associated with the EcR polypeptides of insect ecdysone receptors of the present invention
30 as described in the preceding Examples, are coupled to pins according to the procedure of Geysen *et al.* (1987), and reacted with candidate insecticidal compounds, generally at a

- 68 -

concentration in the range from about 0.05 μ M to about 100 μ M of the candidate compound. The binding of compounds is detected using standard procedures, and compounds having insecticidal activity are identified. Preferably, such compounds exhibit insecticidal activity against a range of insects, including diptera, hemiptera, coleoptera, ants, and moths, amongst others. More preferably, the compounds will exhibit insecticidal activity against *L. cuprina*, *M. persicae*, *D. melanogaster*, scale insect, white fly, and leaf hopper, amongst others. In a particularly preferred embodiment, insecticidal compounds are specific to *L. cuprina* and/or *M. persicae* and close relatives thereof.

10

EXAMPLE 17**Cloned *Myzus persicae* EcR/USP complex binds ponasterone A *in vitro*.**

In vitro-translated *Myzus persicae* EcR (MpEcR) polypeptide and an *in vitro*-translated *M. persicae* USP (MpUSP) polypeptide were produced labelled with [³⁵S]Methionine, using the Promega TNT-Coupled Reticulocyte Lysate System. Each batch of lysate contained 100-200 mg/ml of endogenous proteins (using BSA as a standard). The products were analysed by SDS-PAGE and radioautography. The results confirmed that the cloned cDNAs encode proteins of the sizes predicted from the length of putative open reading frames of the cDNAs present in plasmids pMpEcR and pMpUSP. The yields of EcR and USP were similar as assessed by SDS-PAGE.

20

In functional assays, DNA plasmids pMpEcR (AGAL Accession No. NM99/04567; 1 mg) or pMpUSP (AGAL Accession No. NM99/04568; 1 mg), which have been constructed using the vector pBK-CMV, and 1 ml of appropriate TNT RNA Polymerase were added to 48 ml of reaction mix which contained TNT Lysate, TNT Reaction Buffer, amino acid mixture, Rnasin Ribonuclease Inhibitor and nuclease-free water in volumes specified in the manufacturer's protocol. In control reactions, a Luciferase T3 control DNA (Promega) was used in place of pMpEcR or pMpUSP. T7 RNA Polymerase was used for transcription of the *M. persicae* EcR RNA from plasmid pMpEcR, whilst T3 RNA Polymerase was used for transcription of *M. persicae* USP RNA from the plasmid pMpUSP and the Luciferase T3 control DNA. The reactions were carried out for 90 minutes at 30°C.

30

CONFIDENTIAL

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 69 -

The control reaction produces 150-500 ng of luciferase per 50 ml reaction.

The ecdysteroid binding activities of an *in vitro*-translated *Myzus persicae* EcR (MpEcR) polypeptide and an *in vitro*-translated complex of the *M.persicae* EcR and USP polypeptides were produced from the RNAs using the TNT-Coupled Reticulocyte Lysate System (Promega). The mixtures were stored at -20°C overnight.

After thawing the translation products, 15 ml aliquots of the reaction mixture containing *M.persicae* EcR and USP polypeptides were combined to promote formation of the EcR/USP complex. For assays of individual proteins, 15 ml of the reaction mixture containing *M.persicae* EcR polypeptide or 15 ml of the reaction mixture containing *M.persicae* USP polypeptide was combined with 15 ml of control luciferase protein reaction mixture. Samples were each diluted to 435 ml with EcR40 buffer [40 mM KCl, 25 mM HEPES pH 7.0, 1 mM EDTA, 1mM DTT, BSA(0.5mg/ml), 10% glycerol] to allow for triplicates in the ligand binding assay. A control reaction (Blank) was established which contained EcR40 buffer only. An aliquot (140 ml) of each diluted sample was incubated with tritiated ponasterone A (DuPont NEN, Batch Number 3281108) at a final concentration of 2.2 nM for 90 min at room temperature. After incubation, the ligand binding reactions were placed on ice. The samples were pipetted onto Whatman GF/C filters and incubated for 30 sec. The filters were then placed on a vacuum sinter, washed with 10 ml EcR40 buffer and transferred to scintillation vials. After adding 7 ml of InstaGel Plus to each vial, the contents were vortexed and left at room temperature until the filters became transparent. The receptor bound ligand was quantified using a TriCarb 2100TR scintillation counter.

The results depicted in Figure 4 indicate that significantly higher amounts of ponasterone A bind to the complex than to either the USP or EcR polypeptides alone.

Each reference cited herein is incorporated by reference to the extent that it is not inconsistent with the present disclosure.

30

- 70 -

REFERENCES

1. Amann and Brosius (1985). *Gene* **40**: 183.
2. Ausubel, *et al* (1992), *Current Protocols in Molecular Biology*, Greene/Wiley, New York
3. Bugg, *et al* (1993) *Scientific American*, December Issue: pp 60-66.
- 5 4. Devereux, J., *et al.* (1984). *Nucl. Acids Res.* **12**:387-395.
5. Evans (1988) *Science*, **240**: 889-895.
6. Geysen *et al* (1987) *J. Immunol Methods*, **102**, 259-274.
7. Greene *et al.* (1988) *Nucleic Acids Res.* **15**:369.
8. Hollenberd and Evans (1988) *Cell* **55**: 899-906.
- 10 9. Hollenberg *et al* (1991) *Cell*, **67**, 59-77.
10. Jones, G and Sharp, P. (1997) *Proc. Nat. Acad. Sci. USA* **94**:13499-13503.
11. Kamimura *et al* (1996) *Comp. Biochem. Physiol.* **113**, 341-347.
12. Koelle *et al* (1991) *Cell*, **67**:59-77
13. Krust *et al* (1986) *EMBO. J.*, **5**: 891-897.
- 15 14. Luckow *et al*, (1993) *J. Virol*, **67**: 4566.
15. McBurney *et al* (1991), *Nucleic Acids Res.*, **19**, 5755-5761.
16. McPherson, M.J., Quirke, P. and Taylor, G.R. (1991) *PCR A Practical Approach*. IRL Press, Oxford University Press, Oxford, United Kingdom.
17. Needleman and Wunsch (1970) *J. Mol. Biol.* **48**:443-453.
- 20 18. Potrykus (1990) *Bio/Technology* **8**: 535-542.
19. Riddihough, G., and Pelham, H. R. B., (1987), *EMBO J.*, **6**, 3729-3734
20. Sambrook *et al* (1989), *Molecular Cloning: A Laboratory Manual*, Cold Springs Harbor Laboratory Press
21. Segraves and Hogness (1990) *Genes Devel.* **4**: 204-219.
- 25 22. Shimatake and Rosenberg (1981) *Nature* **292**: 128
23. Studier and Moffat (1986) *J. Mol. Biol.* **182**: 113
24. Thompson, *et al.*, (1994) *Nucl. Acids Res.* **22**:4673-4680.
25. Tzertzinis *et al* (1994) *J. Mol. Biol.* **258**, 479-486.
26. Von Itzstein *et al* (1993) *Nature* Vol. **363**: 418-423.
- 30 27. Yund, *et al.* (1978) *Proc. Nat. Acad. Sci. USA*, **24**: 6039-6043.

667020"02434660

P:\OPER\MRO\CDYSONE.CIP - 1/7/99

- 71 -

WE CLAIM:

1. An isolated nucleic acid molecule comprising a nucleotide sequence which encodes or is complementary to a sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide or a bioactive derivative or analogue thereof, wherein said polypeptide is selected from the groups consisting of (i) an EcR polypeptide of a steroid receptor; (ii) the partner protein (USP polypeptide) of a steroid receptor; and (iii) the USP polypeptide of a juvenile hormone receptor; and wherein said polypeptide comprises an amino acid sequence that is at least 40% identical to an amino acid sequence selected from the group consisting of:
 - (a) an amino acid sequence set forth in any one of <400>2, <400>4, <400>6, <400>10, <400>12 or <400>14; and
 - (b) an amino acid sequence encoded by a cDNA present in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568.
2. The isolated nucleic acid molecule according to claim 1, wherein the steroid receptor is an ecdysteroid receptor.
3. The isolated nucleic acid molecule according to claim 2, wherein the ecdysteroid receptor is an insect ecdysone receptor.
4. The isolated nucleic acid molecule according to claim 3, wherein the insect ecdysone receptor comprises the EcR polypeptide of an insect ecdysone receptor or the partner protein (USP polypeptide) of an insect ecdysone receptor.
5. The isolated nucleic acid molecule according to claim 4, wherein the insect is selected from the list comprising dipteran, hemipteran, coleopteran, lepidopteran, and neuropteran insects and ants.
6. The isolated nucleic acid molecule according to claim 5, wherein the hemipteran insect is *Myzus persicae* or a close relative thereof.

Accepted for publication

P:\QPER\MRO\ECDYSONE.CIP - 1/7/99

- 72 -

7. The isolated nucleic acid molecule according to claim 6, wherein the insect steroid receptor polypeptide comprises an EcR polypeptide of the *M. persicae* ecdysone receptor having the amino acid sequence set forth in <400>6 or <400>10 or encoded by the cDNA present in plasmid pMpEcR (AGAL Accession No. NM99/04567) or a bioactive analogue or derivative thereof.
8. The isolated nucleic acid molecule according to claim 6, wherein the insect steroid receptor polypeptide comprises an EcR partner protein (USP polypeptide) of the *M. persicae* ecdysone receptor or a USP polypeptide of the *M. persicae* juvenile hormone receptor having or including the amino acid sequence set forth in <400>12 or encoded by the cDNA present in plasmid pMpUSP (AGAL Accession No. NM99/04568) or a bioactive analogue or derivative thereof.
9. The isolated nucleic acid molecule according to claim 5, wherein the dipteran insect is *L. cuprina* or a close relative thereof.
10. The isolated nucleic acid molecule according to claim 9, wherein the insect steroid receptor polypeptide comprises an EcR polypeptide of the *L. cuprina* ecdysone receptor having the amino acid sequence set forth in <400>2 or encoded by the cDNA present in plasmid pLcEcR (AGAL Accession No. NM99/04566) or a bioactive analogue or derivative thereof.
11. The isolated nucleic acid molecule according to claim 9, wherein the insect steroid receptor polypeptide comprises an EcR partner protein (USP polypeptide) of the *L. cuprina* ecdysone receptor or a USP polypeptide of the *L. cuprina* juvenile hormone receptor having the amino acid sequence set forth in <400>4 or <400>14 or encoded by the cDNA present in plasmid pLcUSP (AGAL Accession No. NM99/04565) or a bioactive analogue or derivative thereof.
12. The isolated nucleic acid molecule according to claim 1, wherein the bioactive derivative or analogue comprises a fragment of an EcR polypeptide of an insect ecdysone receptor or a fragment of an EcR partner protein (USP polypeptide) of an insect ecdysone

004566024960

P:\OPER\MRO\EC DYSONE.CIP - 1/7/99

- 73 -

receptor, wherein said fragment includes at least one ligand-binding region of said EcR polypeptide or said EcR partner protein (USP polypeptide).

13. The isolated nucleic acid molecule according to claim 12, wherein the ligand-binding region comprises a linker domain of the EcR polypeptide or a linker domain of the EcR partner protein (USP polypeptide).

14. The isolated nucleic acid molecule according to claim 12, wherein the ligand-binding region comprises a hormone-binding domain of the EcR polypeptide or a hormone-binding domain of the EcR partner protein (USP polypeptide).

15. The isolated nucleic acid molecule according to claim 12, wherein the ligand-binding region comprises a linker domain and hormone-binding domain of the EcR polypeptide or a linker domain and hormone-binding domain of the EcR partner protein (USP polypeptide).

16. The isolated nucleic acid molecule according to claim 1, comprising a protein-encoding nucleotide sequence which is at least 40% identical to any one of the nucleotide sequences set forth in <400>1, <400>3, <400>5, <400>9, <400>11 or <400>13 or a complementary nucleotide sequence thereto or the cDNA present in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568.

17. An isolated nucleic acid molecule comprising a nucleotide sequence which encodes or is complementary to a sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide or a bioactive derivative or analogue thereof, wherein said nucleotide sequence is selected from the list comprising:

- (i) a nucleotide sequence having at least 40% identity to any one of the nucleotide sequences set forth in <400>1, <400>3, <400>5, <400>9, <400>11 or <400>13 or a complementary nucleotide sequence thereto;
- (ii) a nucleotide sequence that is capable of hybridising under at least low stringency conditions to any one of the nucleotide sequences set forth in <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11 or <400>13 or to a

667020 " 04/04/99

P:\OTER\MRO\ECDYSONE.CIP - 1/7/99

- 74 -

complementary nucleotide sequence thereto;

(iii) a nucleotide sequence having at least 40% identity to a nucleotide sequence of a cDNA present in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568;

(iv) a nucleotide sequence that is capable of hybridising under at least low stringency conditions to a cDNA present in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568; and

(v) a nucleotide sequence that is amplifiable by PCR using a nucleic acid primer sequence set forth in any one of <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20.

18. An isolated nucleic acid molecule which encodes an insect steroid receptor polypeptide and comprises the nucleotide sequence set forth in <400>1 or a complementary nucleotide sequence thereto or the nucleotide sequence of the cDNA present in plasmid pLcEcR (AGAL Accession No. NM99/04566) .

19. An isolated nucleic acid molecule which encodes an insect steroid receptor polypeptide or a juvenile hormone receptor polypeptide and comprises the nucleotide sequence set forth in <400>3 or <400>13 or a complementary nucleotide sequence thereto or the nucleotide sequence of the cDNA present in plasmid pLcUSP (AGAL Accession No. NM99/04565).

20. An isolated nucleic acid molecule which encodes an insect steroid receptor polypeptide and comprises the nucleotide sequence set forth in <400>5 or <400>7 or <400>8 or <400>9 or a complementary nucleotide sequence thereto or the nucleotide sequence of the cDNA present in plasmid pMpEcR (AGAL Accession No. NM99/04567).

21. An isolated nucleic acid molecule which encodes an insect steroid receptor polypeptide or a juvenile hormone receptor polypeptide and comprises the nucleotide sequence set forth in <400>11 or a complementary nucleotide sequence thereto or the nucleotide sequence of the cDNA present in plasmid pMpUSP (AGAL Accession No. NM99/04568).

- 75 -

22. A method of identifying an isolated nucleic acid molecule which encodes an insect steroid receptor polypeptide which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide comprising:

- (i) hybridising genomic DNA, mRNA or cDNA with a hybridisation-effective amount of one or more probes selected from the list comprising:
 - (a) probes comprising at least 10 contiguous nucleotides in length derived from any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, <400>13, <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20 or a complementary nucleotide sequence thereto;
 - (b) probes comprising at least 10 contiguous nucleotides in length derived from a cDNA contained in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568; and
 - (c) hybridisation probes comprising the nucleotide sequences set forth in any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, or <400>13 or a complementary nucleotide sequence thereto or a homologue, analogue or derivative thereof which is at least 40% identical to said sequence or complement; and
- (ii) detecting the hybridisation.

23. The method of claim 22 wherein the step of detecting the hybridisation comprises detecting a reporter molecule that is covalently bound to the probe.

24. A method of identifying an isolated nucleic acid molecule which encodes an insect steroid receptor polypeptide which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide comprising:

- (i) annealing to genomic DNA, mRNA or cDNA, one or more PCR primers comprising at least 10 contiguous nucleotides in length derived from the group consisting of:
 - (a) a primer derived from any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, <400>13, <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20 or a complementary nucleotide sequence thereto; and

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 76 -

- (b) a primer derived from a cDNA contained in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568; and
- (ii) amplifying a nucleotide sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide in a polymerase chain reaction.

25. A method of identifying an isolated nucleic acid molecule which encodes an insect steroid receptor polypeptide which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide comprising:

- (i) amplifying a nucleotide sequence which encodes a steroid receptor polypeptide or a juvenile hormone receptor polypeptide in a polymerase chain reaction using one or more PCR primers comprising at least 10 contiguous nucleotides in length from the group consisting of:
 - (a) a primer derived from any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, <400>13, <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20 or a complementary nucleotide sequence thereto; and
 - (b) a primer derived from a cDNA contained in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568;
- (ii) hybridising the amplified nucleotide sequence to genomic DNA, mRNA or cDNA with a hybridisation-effective amount of one or more probes selected from the group consisting of:
 - (a) a probe comprising at least 10 contiguous nucleotides in length derived from any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, <400>13, <400>15, <400>16, <400>17, <400>18, <400>19 or <400>20 or a complementary nucleotide sequence thereto;
 - (b) a probe comprising at least 10 contiguous nucleotides in length derived from a cDNA contained in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568; and
 - (c) hybridisation probes comprising the nucleotide sequences set forth in any one of <400>1, <400>3, <400>5, <400>7, <400>8, <400>9, <400>11, or

Accepted for publication

- 77 -

<400>13 or a complementary nucleotide sequence thereto or a homologue, analogue or derivative thereof which is at least 40% identical to said sequence or complement; and

(iii) detecting the hybridisation.

26. The method of claim 25 wherein the step of detecting the hybridisation comprises detecting a reporter molecule that is covalently bound to the probe.

27. The method according to claim 22, further comprising the step of isolating the identified nucleic acid molecule.

28. A genetic construct comprising the isolated nucleic acid molecule according to claim 1 operably linked to a promoter sequence.

29. The genetic construct according to claim 28, wherein the promoter is the SV40, MMTV, polyhedron or p10 promoter.

30. A recombinant or isolated polypeptide comprising a steroid receptor polypeptide or juvenile hormone receptor polypeptide or a bioactive derivative or analogue thereof, wherein said polypeptide:

(i) is selected from the list comprising EcR polypeptide of a steroid receptor, the partner protein (USP polypeptide) of a steroid receptor and the USP polypeptide of a juvenile hormone receptor; and

(ii) comprises an amino acid sequence that is at least 40% identical to any one of the amino acid sequences selected from the group consisting of:

(a) an amino acid sequence set forth in <400>2, <400>4, <400>6, <400>10, <400>12 or <400>14; and

(b) an amino acid sequence encoded by a cDNA present in any one of the plasmids deposited under AGAL Accession Nos. NM99/04565, NM99/04566, NM99/04567, or NM99/04568;

wherein said polypeptide is substantially free of naturally-associated cellular components.

Accepted for publication

- 78 -

31. A cell comprising the nucleic acid molecule according to claim 1.
32. A cell comprising the genetic construct according to claim 28.
33. A cell which expresses the isolated or recombinant polypeptide according to claim 30.
34. A method of identifying a modulator of steroid receptor-mediated gene expression or juvenile hormone receptor-mediated gene expression comprising:
- (i) assaying the expression of a reporter gene in the presence of the recombinant or isolated polypeptide according to claim 30 and a potential modulator; and
 - (ii) assaying the expression of the reporter gene in the presence of the recombinant or isolated polypeptide according to claim 30 and without said potential modulator; and
 - (ii) comparing expression of the reporter gene in the presence of the potential modulator to the expression of a reporter gene in the absence of the potential modulator,
- wherein said reporter gene is placed operably under the control of a steroid response element (SRE) to which said steroid receptor binds or a promoter sequence comprising said SRE.
35. A method of identifying a potential insecticidal compound comprising:
- (i) assaying the binding directly or indirectly of the recombinant or isolated polypeptide according to claim 30 to a steroid response element (SRE) to which said polypeptide binds, in the presence of a candidate compound; and
 - (ii) assaying the binding directly or indirectly of the recombinant or isolated polypeptide according to claim 30 to a steroid response element (SRE) to which said polypeptide binds, in the absence of said candidate compound; and
 - (ii) comparing the binding assayed at (i) and (ii), wherein a difference in the level of binding indicates that the candidate compound possesses potential insecticidal activity.
36. A method of identifying a candidate insecticidally-active agent comprising the steps of:

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 79 -

- a) expressing an EcR polypeptide of an insect steroid receptor or a fragment thereof which includes the ligand-binding region, optionally in association with an EcR partner protein (USP polypeptide) of an insect steroid receptor or ligand binding domain thereof, optionally in association with an insect steroid or analogue thereof so as to form a complex;
- b) purifying or precipitating the complex;
- c) determining the three-dimensional structure of the ligand binding domain of the complex; and
- d) identifying compounds which bind to or associate with the three-dimensional structure of the ligand binding domain, wherein said compounds represent candidate insecticidally-active agents.
37. A method of identifying a candidate insecticidally-active agent comprising the steps of:
- a) expressing a USP polypeptide of a juvenile hormone receptor or a fragment thereof which includes the ligand-binding region, optionally in association with an EcR polypeptide of an insect steroid receptor or ligand binding domain thereof, and optionally in association with an insect steroid or analogue thereof, so as to form a complex;
- b) purifying or precipitating the complex;
- c) determining the three-dimensional structure of the ligand binding domain of the complex; and
- d) identifying compounds which bind to or associate with the three-dimensional structure of the ligand binding domain, wherein said compounds represent candidate insecticidally-active agents.
38. A synthetic compound which interacts with the three dimensional structure of a polypeptide or protein selected from the list comprising:
- (i) an EcR polypeptide of a steroid receptor or a fragment or bioactive derivative thereof;
- (ii) an EcR partner protein (USP polypeptide) of a steroid receptor or a fragment or bioactive derivative thereof;

- 80 -

- (iii) a USP polypeptide of a juvenile hormone receptor or a fragment or bioactive derivative thereof; and
- (iv) a functional receptor or protein complex formed by association of (i) and (ii),
- wherein said compound is capable of binding to said polypeptide or protein to agonise or antagonise the binding activity or bioactivity thereof.

39. A method of identifying a synthetic compound for insecticidal activity comprising contacting the recombinant or isolated polypeptide according to claim 30 with said compound for a time and under conditions sufficient for binding to occur and detecting said binding using a detection means, wherein the occurrence of binding is indicative of potential insecticidal activity of the compound.

[illegible]

- 81 -

ABSTRACT

The present invention provides isolated nucleic acid molecules encoding polypeptides
5 comprising functional steroid hormone and juvenile hormone receptors, in particular isolated
nucleic acid molecules which encode polypeptides comprising the *Lucilia cuprina* and *Myzus*
persicae ecdysone receptors and juvenile hormone receptors. The present invention further
provides functional recombinant steroid and juvenile hormone receptors and recombinant
polypeptide subunits thereof and derivatives and analogues thereof. The present invention
10 further provides screening systems and methods of identifying insecticidally-active agents
which are capable of agonising or antagonising insect receptor function, or alternatively or in
addition, which modify the affinity of said receptors for their cellular stimuli (eg. insect steroids
or juvenile hormones) or analogues thereof, or alternatively or in addition, which act as
insecticides by virtue of their ability to agonise or antagonise the activity of insect hormones.

15

657020 "0434660

651020"02494e60




FIGURE 1

657070" 0279he60




FIGURE 2

3/4





FIGURE 3

4/4

 **^3H -PonA binding by Myzus
in vitro translation products****FIGURE 4**

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 1 -

SEQUENCE LISTING

<110> COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANISATION

<120> NOVEL GENETIC SEQUENCES ENCODING STEROID AND JUVENILE HORMONE

5 RECEPTOR POLYPEPTIDES AND INSECTICIDAL MODALITIES THEREFOR II

```
<130> p:\oper\mro\ecdysone.cip
```

<140> US continuation-in-part of PCT/AU99/00033

<141> 1999-07-02

10 <150> PCT/AU/00033

<151> 1999-01-15

<150> AU PP1536

<151> 1998-01-15

15

<160> 20

<170> PatentIn Ver. 2.0

<210> 1

20 <211> 2274

<212> DNA.

<213> Lucilia cuprina

<220>

25 <221> CDS

<222> (1) . . (2271)

<400> 1

atg atg aaa cga cgt tgg tct aat aat ggc ggt ttt gcc gcc tta aaa 48

30 Met Met Lys Arg Arg Trp Ser Asn Asn Gly Gly Phe Ala Ala Leu Lys

1

5

10

15

- 2 -

30

45

60

80

25

110

125

140

- 3 -

	ggc ctg ggt ggt ggt ggt ggc agt caa gtg aat aat cac aat cac agc	480
	Gly Leu Gly Gly Gly Gly Gly Ser Gln Val Asn Asn His Asn His Ser	
	145 150 155 160	
5	cac aat cat tta cac cac aac agc aac agt aat cac agt aat agc agt	528
	His Asn His Leu His His Asn Ser Asn Ser Asn His Ser Asn Ser Ser	
	165 170 175	
	tcc cac cac aca aat ggc cac atg ggt att ggc ggc ggt ggt ggt ggc	576
10	Ser His His Thr Asn Gly His Met Gly Ile Gly Gly Gly Gly Gly Gly	
	180 185 190	
	tta tcg gtc aat att aat ggt ccc aat atc gtt agc aat gcc caa cag	624
	Leu Ser Val Asn Ile Asn Gly Pro Asn Ile Val Ser Asn Ala Gln Gln	
15	195 200 205	
	tta aac tcg tta cag gcc tca caa aat ggc caa gtt att cat gcc aat	672
	Leu Asn Ser Leu Gln Ala Ser Gln Asn Gly Gln Val Ile His Ala Asn	
	210 215 220	
20	att ggc att cac agt atc atc agt aat gga tta aat cat cat cac cat	720
	Ile Gly Ile His Ser Ile Ile Ser Asn Gly Leu Asn His His His His	
	225 230 235 240	
25	cat cat atg aat aac agt agt atg atg cat cat aca ccc aga tct gaa	768
	His His Met Asn Asn Ser Ser Met Met His His Thr Pro Arg Ser Glu	
	245 250 255	
	tca gct aat tcc ata tca tca ggt cgt gat gat ctt tca ccc tcg agc	816
30	Ser Ala Asn Ser Ile Ser Ser Gly Arg Asp Asp Leu Ser Pro Ser Ser	
	260 265 270	

- 4 -

gaa tgt gtg gtg ccc gaa aac cag tgt gca atg aaa cga cgc gaa aag 1152
Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg Glu Lys
370 375 380

- 6 -

525

540

560

575

590

605

620

640

- 7 -

caa tcg gat aat gca tag 2274

30 Gln Ser Asp Asn Ala

755

- 8 -

115 120 125

[illegible]

- 9 -

140

160

175

190

205

220

240

255

270

285

Lys Lys Gly Pro Ala Pro Arg Leu Gln Glu Glu Leu Cys Leu Val Cys

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 10 -

	290	295	300
	Gly Asp Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly		
	305	310	315 320
5	Cys Lys Gly Phe Phe Arg Arg Ser Val Thr Lys Asn Ala Val Tyr Cys		
	325	330	335
	Cys Lys Phe Gly His Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys		
10	340	345	350
	Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro		
	355	360	365
15	Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg Glu Lys		
	370	375	380
	Lys Ala Gln Lys Glu Lys Asp Lys Ile Gln Thr Ser Val Cys Ala Thr		
	385	390	395 400
20	Glu Ile Lys Lys Glu Ile Leu Asp Leu Met Thr Cys Glu Pro Pro Ser		
	405	410	415
	His Pro Thr Cys Pro Leu Leu Pro Glu Asp Ile Leu Ala Lys Cys Gln		
25	420	425	430
	Ala Arg Asn Ile Pro Pro Leu Ser Tyr Asn Gln Leu Ala Val Ile Tyr		
	435	440	445
30	Lys Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Glu Glu Asp		
	450	455	460

- 11 -

Gln Ser Tyr Tyr Ile Asp Thr Leu Arg Ile Tyr Ile Leu Asn Arg His
30 610 615 620

- 12 -

<213> Lucillia cuprina

P:\OPER\MRO\ECDYSONE,C\P - 1/7/99

- 13 -

<220>

<221> CDS

<222> {1} .. (1401)

5

<400> 1
atg gat aac ggc gag daa gat gct ggg ttc cga ttg gca ccg atg tct 48

10 Met Asp Asn Gly Glu Gln Asp Ala Gly Phe Arg Leu Ala Pro Met Ser
1 5 10 15

15 ccg cag gag ata aag cca gac att tca cta ctc aat gaa aat aat acg 96
Pro Gln Glu Ile Lys Pro **Asp** Ile Ser Leu Leu Asn Glu Asn **Asn** Thr

20 25 30

25 agt agt tat cgg ccc aaa cct gga agt cct aat cca ttt gcc atc gga 144
Ser Ser Tyr Ser Pro Lys Pro Gly Ser Pro Asn Pro Phe Ala Ile Gly
35 40 45

30 ttg cag gca ata aat gca gtc gct gcc gcg aat gcc aat aac caa aat 192
Leu Gln Ala Ile Asn Ala Val Ala Ala Ala Asn Ala Asn Asn Gln Asn

35 50 55 60

caa atg ttg caa act acg cca cca caa cag cag cag tat cca cca aat 240
Gln Met Leu Gln Thr Thr Pro Pro Gln Gln Gln Gln Tyr Pro Pro Asn
65 70 75 80

45 cac ccc ctt agt ggt tgg aaa cac ttg tgt tcc att tgt gga gac cgc 288
50 His Pro Leu Ser Gly Ser Lys His Leu Cys Ser Ile Cys Gly Asp Arg
 85 90 95

55 gcc agt gga aaa cat tat ggg gtc tac agt tgt gag ggt tgt aaa ggg 336
Ala Ser Gly Lys His Tyr Gly Val Tyr Ser Cys Glu Gly Cys Lys Gly
100 105 110

65 ttc ttc aaa cgt acc gta cgc aag gac ttg aca tat gct tgt cgt gag 384
Phe Phe Lys Arg Thr Val Arg Lys Asp Leu Thr Tyr Ala Cys Arg Glu
 115 120 125

70 gac aga aat tgc att ata gat aaa cga caa aga aat cgt tgc cag tat 432
Asp Arg Asn Cys Ile Ile Asp Lys Arg Gln Arg Asn Arg Cys Gln Tyr
75 130 135 140

80 tgt cgt tat caa aag tgt tta gct tgt ggc atg aaa cgc gaa gcg gtc 480
Cys Arg Tyr Gln Lys Cys Leu Ala Cys Gly Met Lys Arg Glu Ala Val

P:OPIRIMROIECDYSONE,CIP - 1/7/99

- 14 -

	145	150	155	160	
5	caa gag gaa cga caa cgt ggt act cgt gct gct aac gct aga gct gct				528
	Gln Glu Glu Arg Gln Arg Gly Thr Arg Ala Ala Asn Ala Arg Ala Ala				
10		165	170	175	
	ggt gct ggc ggt ggt gga gga ggt ggt ggt ggg gta agc aat gtg gtt				576
15	Gly Ala Gly Gly Gly Gly Gly Gly Gly Gly Gly Val Ser Asn Val Val				
		180	185	190	
20	ggt gct ggc gga gaa gac ttt aaa ccc agc agt tca tta cgt gat ctc				624
	Gly Ala Gly Gly Glu Asp Phe Lys Pro Ser Ser Ser Leu Arg Asp Leu				
25		195	200	205	
30	act ata gaa cgc atc att gaa gcc gag caa aag gct gaa tct ttg agc				672
	Thr Ile Glu Arg Ile Ile Glu Ala Glu Gln Lys Ala Glu Ser Leu Ser				
		210	215	220	
35	ggt gat aac gtg ttg ccc ttt ttg cgc gtt ggc aac aat tcc atg gta				720
40	Gly Asp Asn Val Leu Pro Phe Leu Arg Val Gly Asn Asn Ser Met Val				
		225	230	235	240
45	caa cac gac tac aaa ggc gcg gta tct cat ctc tgc cag atg gtt aac				768
	Gln His Asp Tyr Lys Gly Ala Val Ser His Leu Cys Gln Met Val Asn				
50		245	250	255	
55	aaa caa ctc tac caa atg gtt gaa tat gca cgt cga aca cca cat ctt				816
	Lys Gln Leu Tyr Gln Met Val Glu Tyr Ala Arg Arg Thr Pro His Phe				
		260	265	270	
60	aca cat ttg cag cgt gag gat cag ata cta ttg tta aag gct ggc tgg				864
	Thr His Leu Gln Arg Glu Asp Gln Ile Leu Leu Leu Lys Ala Gly Trp				
65		275	280	285	
70	aat gaa ctg cta att gca aat gtt gcc tgg tgc agt att gag tct ctg				912
	Asn Glu Leu Leu Ile Ala Asn Val Ala Trp Cys Ser Ile Glu Ser Leu				
		290	295	300	
75	gat gcc gaa cat gcc tct cct ggt acg gta cat gac ggt tct ttt ggt				960
80	Asp Ala Glu Tyr Ala Ser Pro Gly Thr Val His Asp Gly Ser Phe Gly				
		305	310	315	320
85	egg cgt tca cca gtg cgt cag ccc caa caa ctc ttc ctt aat cag aat				1008
	Arg Arg Ser Pro Val Arg Gln Pro Gln Gln Leu Phe Leu Asn Gln Asn				
90		325	330	335	

667020 "04754660

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 15 -

	ttc tgc tat cat cgc aat agt gct att aag gcc aat gtt gtt tca att	1056
	Phe Ser Tyr His Arg Asn Ser Ala Ile Lys Ala Asn Val Val Ser Ile	
5	340 345 350	
10	ttc gat cgt atc ctc tcg gag ttg agc atc aaa atg aaa cgt ctt aac	1104
	Phe Asp Arg Ile Leu Ser Glu Leu Ser Ile Lys Met Lys Arg Leu Asn	
	355 360 365	
15		
20	atc gat cgc tcg gag ttg tcg tgt ctg aag gca atc ata ctc ttc aat	1152
	Ile Asp Arg Ser Glu Leu Ser Cys Leu Lys Ala Ile Ile Leu Phe Asn	
	370 375 380	
25	cca gac ata cgc ggt ctg aaa tgt cga gcc gac gtc gag gta tgt cgt	1200
	Pro Asp Ile Arg Gly Leu Lys Cys Arg Ala Asp Val Glu Val Cys Arg	
30	385 390 395 400	
35	gaa aaa atc tat gcc tgt ctg gac gaa cac tgc cgc aca gaa cat tca	1248
	Glu Lys Ile Tyr Ala Cys Leu Asp Glu His Cys Arg Thr Glu His Pro	
	405 410 415	
40	ggc gat gat ggc cgc ttt gct cag cta cta cta agg ttg ccc gca ttg	1296
	Gly Asp Asp Gly Arg Phe Ala Gln Leu Leu Leu Arg Leu Pro Ala Leu	
45	420 425 430	
50	cgt tcc atc agt ctc aaa tgt ctc gat cat ttg ttt ttc ttc cgt tta	1344
	Arg Ser Ile Ser Leu Lys Cys Leu Asp His Leu Phe Phe Phe Arg Leu	
	435 440 445	
55		
60	ata ggc gaa aga gca ttg gag gaa tta att gct gag caa ctg gaa gct	1392
	Ile Gly Glu Arg Ala Leu Glu Glu Leu Ile Ala Glu Gln Leu Glu Ala	
	450 455 460	
65	cct atc tgc	1401
	Pro Ile Cys	
70	465	
75	<210> 4	
	<211> 467	
80	<212> PRT	
	<213> Lucillia cuprina	
85	<400> 4	
	Met Asp Asn Gly Glu Gln Asp Ala Gly Phe Arg Leu Ala Pro Met Ser	
90	1 5 10 15	

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 16 -

	Pro	Gln	Glu	Ile	Lys	Pro	Asp	Ile	Ser	Leu	Leu	Asn	Glu	Asn	Asn	Thr
5	20				25				30							
10	Ser	Ser	Tyr	Ser	Pro	Lys	Pro	Gly	Ser	Pro	Asn	Pro	Phe	Ala	Ile	Gly
	35				40				45							
15	Leu	Gln	Ala	Ile	Asn	Ala	Val	Ala	Ala	Ala	Asn	Ala	Asn	Asn	Gln	Asn
	50				55				60							
20	Gln	Met	Leu	Gln	Thr	Thr	Pro	Pro	Gln	Gln	Gln	Gln	Tyr	Pro	Pro	Asn
	65				70				75				80			
25	His	Pro	Leu	Ser	Gly	Ser	Lys	His	Leu	Cys	Ser	Ile	Cys	Gly	Asp	Arg
	85				90				95							
30	Ala	Ser	Gly	Lys	His	Tyr	Gly	Val	Tyr	Ser	Cys	Glu	Gly	Cys	Lys	Gly
	100				105				110							
35	Phe	Phe	Lys	Arg	Thr	Val	Arg	Lys	Asp	Leu	Thr	Tyr	Ala	Cys	Arg	Glu
40	115				120				125							
45	Asp	Arg	Asn	Cys	Ile	Ile	Asp	Lys	Arg	Gln	Arg	Asn	Arg	Cys	Gln	Tyr
	130				135				140							
50	Cys	Arg	Tyr	Gln	Lys	Cys	Leu	Ala	Cys	Gly	Met	Lys	Arg	Glu	Ala	Val
	145				150				155				160			
55	Gln	Glu	Glu	Arg	Gln	Arg	Gly	Thr	Arg	Ala	Ala	Asn	Ala	Arg	Ala	Ala
	165				170				175							
60	Gly	Ala	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Val	Ser	Asn	Val	Val
	180				185				190							
65	Gly	Ala	Gly	Gly	Glu	Asp	Phe	Lys	Pro	Ser	Ser	Ser	Leu	Arg	Asp	Leu
70	195				200				205							
75	Thr	Ile	Glu	Arg	Ile	Ile	Glu	Ala	Glu	Gln	Lys	Ala	Glu	Ser	Leu	Ser
	210				215				220							
80	Gly	Asp	Asn	Val	Leu	Pro	Phe	Leu	Arg	Val	Gly	Asn	Asn	Ser	Met	Val
	225				230				235				240			
85	Gln	His	Asp	Tyr	Lys	Gly	Ala	Val	Ser	His	Leu	Cys	Gln	Met	Val	Asn
	245				250				255							
90	Lys	Gln	Leu	Tyr	Gln	Met	Val	Glu	Tyr	Ala	Arg	Arg	Thr	Pro	His	Phe

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 17 -

	260	265	270
5	Thr His Leu Gln Arg Glu Asp Gln Ile Leu Leu Leu Lys Ala Gly Trp 275	280	285
10	Asn Glu Leu Leu Ile Ala Asn Val Ala Trp Cys Ser Ile Glu Ser Leu 290	295	300
15	Asp Ala Glu Tyr Ala Ser Pro Gly Thr Val His Asp Gly Ser Phe Gly 305	310	315 320
20	Arg Arg Ser Pro Val Arg Gln Pro Gln Gln Leu Phe Leu Asn Gln Asn 325	330	335
25	Phe Ser Tyr His Arg Asn Ser Ala Ile Lys Ala Asn Val Val Ser Ile 340	345	350
30	Phe Asp Arg Ile Leu Ser Glu Leu Ser Ile Lys Met Lys Arg Leu Asn 355	360	365
35	Ile Asp Arg Ser Glu Leu Ser Cys Leu Lys Ala Ile Ile Leu Phe Asn 370	375	380
40	Pro Asp Ile Arg Gly Leu Lys Cys Arg Ala Asp Val Glu Val Cys Arg 385	390	395 400
45	Glu Lys Ile Tyr Ala Cys Leu Asp Glu His Cys Arg Thr Glu His Pro 405	410	415
50	Gly Asp Asp Gly Arg Phe Ala Gln Leu Leu Leu Arg Leu Pro Ala Leu 420	425	430
55	Arg Ser Ile Ser Leu Lys Cys Leu Asp His Leu Phe Phe Phe Arg Leu 435	440	445
60	Ile Gly Glu Arg Ala Leu Glu Glu Leu Ile Ala Glu Gln Leu Glu Ala 450	455	460
65	Pro Ile Cys 465		
70			
75			
80			

667020 "02494660

PAOPERMKOECDYSONE.CIP. 1/7/99

- 18 -

<210> 5

<211> 585

<212> DNA

<213> Myzus persicae

5

<220>

<221> CDS

<222> (1) .. (585)

10 <400> 5

gaa ttc ggc acg agc gcc att gtt aat gga ttt atc cgc acc att agt 48

Glu Phe Gly Thr Ser Ala Ile Val Asn Gly Phe Ile Arg Thr Ile Ser

1 5 10 15

15 ttg atc ctt att ttt ctt ctt ctt ttt ctt tgg agg ttg ttg gcc ttc 96

Leu Ile Leu Ile Phe Leu Leu Leu Phe Leu Trp Arg Leu Leu Ala Phe

20 25 30

cgg ttc ttg ttt ata tct gaa caa cca cct ccc gaa gag ctg tgc ctg 144

20 Arg Phe Leu Phe Ile Ser Glu Gln Pro Pro Pro Glu Glu Leu Cys Leu

35 40 45

gtg tgt ggc gac cgg tgg tcc ggt tac cat tac aac gct ctg aca tgc 192

Val Cys Gly Asp Arg Ser Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys

25 50 55 60

gaa gga tgc aag ggg ttc ttc cgg agg agc atc acc aag aac gcc gtg 240

Glu Gly Cys Lys Gly Phe Phe Arg Arg Ser Ile Thr Lys Asn Ala Val

65 70 75 80

30

tac cag tgc aag tac ggc aac aat tgc gaa atc gac atg tac atg agg 288

667020243460

P:\OPER\MRO\EC DYSONE.CIP - 1/7/99

- 19 -

Tyr Gln Cys Lys Tyr Gly Asn Asn Cys Glu Ile Asp Met Tyr Met Arg

85

90

95

egg aag tgc cag gag tgc egg ctg aaa aaa tgc ctg acc gtc ggc atg 336

5 Arg Lys Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu Thr Val Gly Met

100

105

110

agg cct gaa tgc gtt gta cct gaa gtt caa tgc gca gta aaa aga aag 384

Arg Pro Glu Cys Val Val Pro Glu Val Gln Cys Ala Val Lys Arg Lys

10

115

120

125

gag aaa aaa gct caa cga gaa aaa gat aaa cca aat tct act aca gac 432

Glu Lys Lys Ala Gln Arg Glu Lys Asp Lys Pro Asn Ser Thr Thr Asp

130

135

140

15

att tct cct gaa ata ata aaa ata gaa cct aca gag atg aag att gaa 480

Ile Ser Pro Glu Ile Ile Lys Ile Glu Pro Thr Glu Met Lys Ile Glu

145

150

155

160

20

tgt ggt gaa cca atg ata atg ggc aca cct atg ccg act gta cct tac 528

Cys Gly Glu Pro Met Ile Met Gly Thr Pro Met Pro Thr Val Pro Tyr

165

170

175

gtg aaa cct ttg agt tct ctc gtg ccg aat tcg gca cga gtc acg ggt 576

25 Val Lys Pro Leu Ser Ser Leu Val Pro Asn Ser Ala Arg Val Thr Gly

180

185

190

tac aaa ttt

585

Tyr Lys Phe

30

195

P:IOPERIMROIECDYSONE.CIP.1/7/99

- 20 -

<210> 6

<211> 195

<212> PRT

5 <213> Myzus persicae

<400> 6

Glu Phe Gly Thr Ser Ala Ile Val Asn Gly Phe Ile Arg Thr Ile Ser

1

5

10

15

10

Leu Ile Leu Ile Phe Leu Leu Leu Phe Leu Trp Arg Leu Leu Ala Phe

20

25

30

Arg Phe Leu Phe Ile Ser Glu Gln Pro Pro Pro Glu Glu Leu Cys Leu

15

35

40

45

Val Cys Gly Asp Arg Ser Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys

50

55

60

20 Glu Gly Cys Lys Gly Phe Phe Arg Arg Ser Ile Thr Lys Asn Ala Val

65

70

75

80

Tyr Gln Cys Lys Tyr Gly Asn Asn Cys Glu Ile Asp Met Tyr Met Arg

85

90

95

25

Arg Lys Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu Thr Val Gly Met

100

105

110

Arg Pro Glu Cys Val Val Pro Glu Val Gln Cys Ala Val Lys Arg Lys

30

115

120

125

667000445460

PAOPKIMROECDOYSONE.CIP-1/7/99

- 21 -

Glu Lys Lys Ala Gln Arg Glu Lys Asp Lys Pro Asn Ser Thr Thr Asp

130

135

140

Ile Ser Pro Glu Ile Ile Lys Ile Glu Pro Thr Glu Met Lys Ile Glu

5

145

150

155

160

Cys Gly Glu Pro Met Ile Met Gly Thr Pro Met Pro Thr Val Pro Tyr

165

170

175

10 Val Lys Pro Leu Ser Ser Leu Val Pro Asn Ser Ala Arg Val Thr Gly

180

185

190

Tyr Lys Phe

195

15

<210> 7

<211> 208

<212> DNA

<213> Myzus persicae

20

<400> 7

catgcctgca ggtcgactct agaggatccc ctggtccggt taccattaca acgcactcac 60

ctgtgaaggc tgtaaggggt tctttcgacg gagtgttacc aaaaatgcgg tgtattgttg 120

25

taaatttggt catgcctgcg aaatggacat gtatatgcga cgtaaagtgc aggaatgtag 180

gctgaaaaaa tgtttggtg tgggcatg

208

30

<210> 8

<211> 436

<212> DNA

35

<213> Myzus persicae

<400> 8

catgcggcgg gaatgtgtgg tgcccgaaaa ccagtgtgca atgaaacgac gcgaaaagaa 60

667020"04hshE60

- 22 -

[illegible]

- 23 -

	Pro	Gln	Gln	Gln	Val	Pro	Pro	Ser	Arg	Asn	Gly	Cys	Ser	Thr	Leu	Phe	
					85						90					95	
5	agc	gac	atc	gct	ggc	gtc	aag	cga	ctc	agg	ccc	gac	gat	tgg	ttg	gcc	336
10	Ser	Asp	Ile	Ala	Gly	Val	Lys	Arg	Leu	Arg	Pro	Asp	Asp	Trp	Leu	Ala	
				100					105					110			
15	gtc	aac	tgc	ccg	ccc	gcc	tct	tgc	ccc	ggc	acg	tgc	cac	ata	tcc	tac	384
	Val	Asn	Ser	Pro	Pro	Ala	Ser	Ser	Pro	Gly	Thr	Ser	His	Ile	Ser	Tyr	
20				115					120					125			
25	aca	gtc	ata	tgc	aac	ggc	ggc	ggc	ggc	ggc	ggc	ggc	ggc	ggc	ggc	ggc	432
	Thr	Val	Ile	Ser	Asn	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	
				130				135						140			
30	tac	aac	acg	tct	cca	atg	tgc	acc	aac	agc	tac	gac	ccg	tac	agt	ccg	480
	Tyr	Asn	Thr	Ser	Pro	Met	Ser	Thr	Asn	Ser	Tyr	Asp	Pro	Tyr	Ser	Pro	
35						145			150				155			160	
40	atg	agt	gga	aaa	atc	gtc	aaa	gaa	gag	ttg	tct	ccg	cca	aac	agc	ctg	528
	Met	Ser	Gly	Lys	Ile	Val	Lys	Glu	Glu	Leu	Ser	Pro	Pro	Asn	Ser	Leu	
					165				170						175		
45	tgc	gga	gtc	agc	agc	cat	tgc	gat	ggg	ttg	aag	aag	aag	aaa	ctc	aac	576
50	Ser	Gly	Val	Ser	Ser	His	Ser	Asp	Gly	Leu	Lys	Lys	Lys	Lys	Leu	Asn	
				180					185					190			
55	cac	acg	ccc	tgc	acc	ggc	gtc	gtc	aac	acc	tgc	gca	tgc	ggc	ccc	ggg	624
	His	Thr	Pro	Ser	Thr	Gly	Val	Val	Asn	Thr	Ser	Ala	Ser	Gly	Pro	Gly	
60				195					200					205			
65	ggc	ggc	gtt	ggc	ggc	aat	gtg	ctg	aac	aac	cga	cct	ccc	gaa	gag	ctg	672
	Gly	Gly	Val	Gly	Gly	Asn	Val	Leu	Asn	Asn	Arg	Pro	Pro	Glu	Glu	Leu	
			210					215					220				
70	tgc	ctg	gtg	tgt	ggc	gac	cgg	tgc	tcc	ggc	tac	cat	tac	aac	gct	ctc	720
	Cys	Leu	Val	Cys	Gly	Asp	Arg	Ser	Ser	Gly	Tyr	His	Tyr	Asn	Ala	Leu	
75				225			230				235					240	
80	aca	tgc	gaa	gga	tgc	aag	ggg	ttc	ttc	cgg	agg	agc	atc	acc	aag	aac	768
	Thr	Cys	Glu	Gly	Cys	Lys	Gly	Phe	Phe	Arg	Arg	Ser	Ile	Thr	Lys	Asn	
					245					250					255		
85	gcc	gtg	tac	cag	tgc	aag	tac	ggc	aac	aat	tgc	gaa	atc	gac	atg	tac	816
90	Ala	Val	Tyr	Gln	Cys	Lys	Tyr	Gly	Asn	Asn	Cys	Glu	Ile	Asp	Met	Tyr	
				260					265					270			

- 24 -

	atg	agg	cgg	aag	tgc	cag	gag	tgc	cgg	ctg	aaa	aaa	tgc	ctg	acc	gtc	864
5	Met	Arg	Arg	Lys	Cys	Gln	Glu	Cys	Arg	Leu	Lys	Lys	Cys	Leu	Thr	Val	
			275					280					285				
10	ggc	atg	agg	cct	gaa	tgt	gtt	gta	cct	gaa	gtt	caa	tgc	gca	gta	aaa	912
	Gly	Met	Arg	Pro	Glu	Cys	Val	Val	Pro	Glu	Val	Gln	Cys	Ala	Val	Lys	
15			290				295					300					
20	aga	aag	gag	aaa	aaa	gct	caa	cga	gaa	aaa	gat	aaa	cca	aat	tct	act	960
	Arg	Lys	Glu	Lys	Lys	Ala	Gln	Arg	Glu	Lys	Asp	Lys	Pro	Asn	Ser	Thr	
	305					310					315				320		
25	aca	gac	att	tct	cct	gaa	ata	ata	aaa	ata	gaa	cct	aca	gag	atg	aag	1008
30	Thr	Asp	Ile	Ser	Pro	Glu	Ile	Ile	Lys	Ile	Glu	Pro	Thr	Glu	Met	Lys	
					325					330					335		
35	att	gaa	tgt	ggg	gaa	cca	atg	ata	atg	ggc	aca	cct	atg	ccg	act	gta	1056
	Ile	Glu	Cys	Gly	Glu	Pro	Met	Ile	Met	Gly	Thr	Pro	Met	Pro	Thr	Val	
40				340					345				350				
45	cct	tac	gtg	aaa	cct	ctg	agt	tct	gaa	caa	aaa	gaa	ctg	atc	cac	cga	1104
	Pro	Tyr	Val	Lys	Pro	Leu	Ser	Ser	Glu	Gln	Lys	Glu	Leu	Ile	His	Arg	
			355					360					365				
50	ctt	gtc	tat	ttc	cag	gat	caa	tac	gaa	gct	cct	agt	gaa	aaa	gac	atg	1152
	Leu	Val	Tyr	Phe	Gln	Asp	Gln	Tyr	Glu	Ala	Pro	Ser	Glu	Lys	Asp	Met	
55			370				375					380					
60	aaa	cgt	tta	aca	ata	aat	aat	caa	aac	atg	gat	gaa	tac	gat	gaa	gaa	1200
	Lys	Arg	Leu	Thr	Ile	Asn	Asn	Gln	Asn	Met	Asp	Glu	Tyr	Asp	Glu	Glu	
	385				390					395					400		
65	aaa	caa	agt	gac	acc	aca	tat	cga	atc	atc	act	gag	atg	aca	ata	ctc	1248
70	Lys	Gln	Ser	Asp	Thr	Thr	Tyr	Arg	Ile	Ile	Thr	Glu	Met	Thr	Ile	Leu	
					405				410					415			
75	aca	gtt	caa	ctg	att	gtt	gag	ttt	gcc	aaa	cga	tta	cca	ggg	ttc	gat	1296
	Thr	Val	Gln	Leu	Ile	Val	Glu	Phe	Ala	Lys	Arg	Leu	Pro	Gly	Phe	Asp	
80				420				425					430				
85	aaa	ctt	gta	aga	gaa	gat	caa	atc	act	tta	ctc	aag	gct	tgc	tca	agt	1344
	Lys	Leu	Val	Arg	Glu	Asp	Gln	Ile	Thr	Leu	Leu	Lys	Ala	Cys	Ser	Ser	
			435				440					445					
90	gaa	gct	atg	atg	ttc	agg	gta	gca	agg	aag	tat	gac	atc	acc	act	gac	1392

PAOPER\MRO\BCDYSONE.CIP - 1/7/99

- 25 -

Glu Ala Met Met Phe Arg Val Ala Arg Lys Tyr Asp Ile Thr Thr Asp
 450 455 460

5

tca ata gtg ttt gct aac aac cag cca ttt tca gct gat tca tat aac 1440

10 Ser Ile Val Phe Ala Asn Asn Gln Pro Phe Ser Ala Asp Ser Tyr Asn
 465 470 475 480

15 aaa gct gga ttg gga gat gcc att gaa aac caa ctg tca ttc agt cgg 1488
 Lys Ala Gly Leu Gly Asp Ala Ile Glu Asn Gln Leu Ser Phe Ser Arg
 485 490 495

20

ttt atg cac aat atg aag gtg gat aac gca gaa tat gcc tta ttg acc 1536

25 Phe Met Tyr Asn Met Lys Val Asp Asn Ala Glu Tyr Ala Leu Leu Thr
 500 505 510

30 gcc atc gtc ata ttt tgg agt agg cca aat tta cta gat ggt tgg aaa 1584
 Ala Ile Val Ile Phe Ser Ser Arg Pro Asn Leu Leu Asp Gly Trp Lys
 515 520 525

35

gtg gag aaa atc caa gaa atc tac cta gag tcc tta aaa gct tat gta 1632
 Val Glu Lys Ile Gln Glu Ile Tyr Leu Glu Ser Leu Lys Ala Tyr Val
 530 535 540

40

gat aat cga gac cgt gac aca gca act gta cga tat gog cga ctt ctc 1680

50 Asp Asn Arg Asp Arg Asp Thr Ala Thr Val Arg Tyr Ala Arg Leu Leu
 545 550 555 560

55 tca gta ctt aca gaa ttg cgc aca tca ggc aat gaa aac tct gag cta 1728
 Ser Val Leu Thr Glu Leu Arg Thr Leu Gly Asn Glu Asn Ser Glu Leu
 565 570 575

60

tgt atg aca ctg aaa ctg aaa aac aga gta gta ccc cca ttc ttg gcc 1776

65 Cys Met Thr Leu Lys Leu Lys Asn Arg Val Val Pro Pro Phe Leu Ala
 580 585 590

70

gaa ata tgg gat gtc atg cca 1797
 Glu Ile Trp Asp Val Met Pro
 595

75

80 <210> 10
 <211> 599

85 <212> PRT
 <213> Myzus persicae

90 <400> 10

BCDYSONE

P:\OPER\MRO\ECDYSONE.CIP - 1/7/99

- 26 -

Met Met Asp Gln Lys Cys Asp Val Gly Gly Gly Gly Val Ala Ala Ala
 1 5 10 15
 5
 Ala Ala Gly Ile Gly Gly Gly Gly Val Gly Gly Leu Met Ser Tyr Asn
 20 25 30
 10
 Arg Gly Arg Gly Gly Thr Glu Val Ile Ile Lys Pro Arg Ser Pro Ala
 35 40 45
 15
 Val Val Gln Val Ala Thr Gly Gly Ser Tyr His Gly Leu Pro Ala Ala
 50 55 60
 20
 Ser Asp Ala Val Ile Val Arg Ser Pro Pro Gly Gly His Leu Pro Gly
 65 70 75 80
 25
 Pro Gln Gln Gln Val Pro Pro Ser Arg Asn Gly Cys Ser Thr Leu Phe
 85 90 95
 30
 Ser Asp Ile Ala Gly Val Lys Arg Leu Arg Pro Asp Asp Trp Leu Ala
 100 105 110
 40
 Val Asn Ser Pro Pro Ala Ser Ser Pro Gly Thr Ser His Ile Ser Tyr
 115 120 125
 45
 Thr Val Ile Ser Asn Gly Gly Gly Gly Gly Gly Gly Gly Gly Gly
 130 135 140
 50
 Tyr Asn Thr Ser Pro Met Ser Thr Asn Ser Tyr Asp Pro Tyr Ser Pro
 145 150 155 160
 55
 Met Ser Gly Lys Ile Val Lys Glu Glu Leu Ser Pro Pro Asn Ser Leu
 165 170 175
 60
 Ser Gly Val Ser Ser His Ser Asp Gly Leu Lys Lys Lys Lys Leu Asn
 180 185 190
 70
 His Thr Pro Ser Thr Gly Val Val Asn Thr Ser Ala Ser Gly Pro Gly
 195 200 205
 75
 Gly Gly Val Gly Gly Asn Val Leu Asn Asn Arg Pro Pro Glu Glu Leu
 210 215 220
 80
 Cys Leu Val Cys Gly Asp Arg Ser Ser Gly Tyr His Tyr Asn Ala Leu
 225 230 235 240
 85
 Thr Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Ser Ile Thr Lys Asn
 90

03
 06
 08
 09
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 00

- 27 -

	245	250	255
5	Ala Val Tyr Gln Cys Lys Tyr Gly Asn Asn Cys Glu Ile Asp Met Tyr 260 265 270		
10	Met Arg Arg Lys Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu Thr Val 275 280 285		
15	Gly Met Arg Pro Glu Cys Val Val Pro Glu Val Gln Cys Ala Val Lys 290 295 300		
20	Arg Lys Glu Lys Lys Ala Gln Arg Glu Lys Asp Lys Pro Asn Ser Thr 305 310 315 320		
25	Thr Asp Ile Ser Pro Glu Ile Ile Lys Ile Glu Pro Thr Glu Met Lys 325 330 335		
30	Ile Glu Cys Gly Glu Pro Met Ile Met Gly Thr Pro Met Pro Thr Val 340 345 350		
35	Pro Tyr Val Lys Pro Leu Ser Ser Glu Gln Lys Glu Leu Ile His Arg 355 360 365		
40	Leu Val Tyr Phe Gln Asp Gln Tyr Glu Ala Pro Ser Glu Lys Asp Met 370 375 380		
45	Lys Arg Leu Thr Ile Asn Asn Gln Asn Met Asp Glu Tyr Asp Glu Glu 385 390 395 400		
50	Lys Gln Ser Asp Thr Thr Tyr Arg Ile Ile Thr Glu Met Thr Ile Leu 405 410 415		
55	Thr Val Gln Leu Ile Val Glu Phe Ala Lys Arg Leu Pro Gly Phe Asp 420 425 430		
60	Lys Leu Val Arg Glu Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser 435 440 445		
65	Glu Ala Met Met Phe Arg Val Ala Arg Lys Tyr Asp Ile Thr Thr Asp 450 455 460		
70	Ser Ile Val Phe Ala Asn Asn Gln Pro Phe Ser Ala Asp Ser Tyr Asn 465 470 475 480		
75	Lys Ala Gly Leu Gly Asp Ala Ile Glu Asn Gln Leu Ser Phe Ser Arg 485 490 495		

P:\OPERIMRO\ECDYSONE.CIF - 1/7/99

- 28 -

5 Phe Met Tyr Asn Met Lys Val Asp Asn Ala Glu Tyr Ala Leu Leu Thr
 500 505 510

10 Ala Ile Val Ile Phe Ser Ser Arg Pro Asn Leu Leu Asp Gly Trp Lys
 515 520 525

15 Val Glu Lys Ile Gln Glu Ile Tyr Leu Glu Ser Leu Lys Ala Tyr Val
 530 535 540

20 Asp Asn Arg Asp Arg Asp Thr Ala Thr Val Arg Tyr Ala Arg Leu Leu
 545 550 555 560

25 Ser Val Leu Thr Glu Leu Arg Thr Leu Gly Asn Glu Asn Ser Glu Leu
 565 570 575

30 Cys Met Thr Leu Lys Leu Lys Asn Arg Val Val Pro Pro Phe Leu Ala
 580 585 590

35 Glu Ile Trp Asp Val Met Pro
 595

40

45 <210> 11
 <211> 1131
 50 <212> DNA
 <213> Myzus persicae
 55 <220>
 <221> CDS
 60 <222> (1)..(1131)
 <400> 11

65 atg tat tcc aac tcg tac acc atg tat tca agt gac aga tta tac agc 48
 Met Tyr Ser Asn Ser Tyr Thr Met Tyr Ser Ser Asp Arg Leu Tyr Ser
 1 5 10 15

70 gtc gat cgg aac agt atg atg aat aat tct tgc aac gta caa gac tct 96
 Val Asp Arg Asn Ser Met Met Asn Asn Ser Cys Asn Val Gln Asp Ser
 20 25 30

75 ccg aat tac ccg ccc aac cat cca ctc agc ggt tcg aaa cat ctg tgc 144
 Pro Asn Tyr Pro Pro Asn His Pro Leu Ser Gly Ser Lys His Leu Cys
 85 35 40 45

90 tcc ata tgc ggc gat cgc gcc agt gga aaa cat tac gga gtc tac agc 192
 Ser Ile Cys Gly Asp Arg Ala Ser Gly Lys His Tyr Gly Val Tyr Ser

66700343660

P:\OPER\MRO\EC\DYSONE.CIP - 1/7/99

- 29 -

	50	55	60	
5	tgc gag ggg tgc aaa ggg ttc ttc aaa cgc aca gtg agg aaa aat ttg			240
	Cys Glu Gly Cys Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asn Leu			
10	65	70	75	80
	tca tac gcg tgt cgc gaa gaa aac aaa tgc atc atc gac aag cgc caa			288
15	Ser Tyr Ala Cys Arg Glu Glu Asn Lys Cys Ile Ile Asp Lys Arg Gln			
	85	90	95	
20	cga aat cgg tgc caa tac tgc agg tat caa aaa tgt ttg acc atg ggc			336
	Arg Asn Arg Cys Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Thr Met Gly			
25	100	105	110	
	atg aaa aga gaa gct gtg cag gaa gaa aga caa cgt aca aaa gaa cga			384
30	Met Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Thr Lys Glu Arg			
	115	120	125	
35	gat cat aat aac atc gaa gtt gaa ccc acg agc agt tct aat act gat			432
40	Asp His Asn Asn Ile Glu Val Glu Pro Thr Ser Ser Ser Asn Thr Asp			
	130	135	140	
45	atg cca gtg gaa ctc ata tta agg gct gag aat aaa gct gat gct ata			480
	Met Pro Val Glu Leu Ile Leu Arg Ala Glu Asn Lys Ala Asp Ala Ile			
50	145	150	155	160
	aag act gaa caa cag tat ata gag caa cga cat cct caa cat act gtt			528
55	Lys Thr Glu Gln Gln Tyr Ile Glu Gln Arg His Pro Gln His Thr Val			
	165	170	175	
60	ggc gct att tgt caa gca act gac aag cag tta ata caa ctt gtc gaa			576
	Gly Ala Ile Cys Gln Ala Thr Asp Lys Gln Leu Ile Gln Leu Val Glu			
65	180	185	190	
	tgg gcc aag cat ata ccg cat ttt aaa aat tta cct cta ggc gat caa			624
70	Trp Ala Lys His Ile Pro His Phe Lys Asn Leu Pro Leu Gly Asp Gln			
	195	200	205	
75	gtt tta tta ttg aga gct ggt tgg aat gag ttg atg att gca gca ttt			672
80	Val Leu Leu Leu Arg Ala Gly Trp Asn Glu Leu Met Ile Ala Ala Phe			
	210	215	220	
85	tcc cat aga tca atc agt gta aaa gat ggt ata gtc tta gct act gga			720
	Ser His Arg Ser Ile Ser Val Lys Asp Gly Ile Val Leu Ala Thr Gly			
90	225	230	235	240

CC-BY-NC-SA 4.0 International

- 30 -

	ctt act gtt gac aga gat tca gct cac caa gct ggt gtt gaa gct ata	768
5	Leu Thr Val Asp Arg Asp Ser Ala His Gln Ala Gly Val Glu Ala Ile	
	245 250 255	
10	ttt gat cgt gta ctc act gaa ctc gtt gct aaa atg aga gat atg ggt	816
	Phe Asp Arg Val Leu Thr Glu Leu Val Ala Lys Met Arg Asp Met Gly	
	260 265 270	
15	atg gat aga aca gag ctt ggc tgt ttg cgt act att att ctt ttt aat	864
20	Met Asp Arg Thr Glu Leu Gly Cys Leu Arg Thr Ile Ile Leu Phe Asn	
	275 280 285	
25	cca ggt tca aaa ggt ttg cag tct gtg aat gaa gtg caa gta ctg cgt	912
	Pro Gly Ser Lys Gly Leu Gln Ser Val Asn Glu Val Gln Val Leu Arg	
30	290 295 300	
35	gat aag gtt tat gtt gcg tta gaa gaa tat tgt cgt aca aca cat cca	960
	Asp Lys Val Tyr Val Ala Leu Glu Glu Tyr Cys Arg Thr Thr His Pro	
	305 310 315 320	
40	gaa gaa cct gga cga ttt gct aaa cta ctt ctt cgg ctt cct tca tta	1008
	Glu Glu Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ser Leu	
45	325 330 335	
50	cgt tca att gga tta aaa tgt ctg gaa cat tta ttc ttt tat aaa ctt	1056
	Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Tyr Lys Leu	
	340 345 350	
55	att ggc gat tcc cca att gat aca ttt tta atg gaa gtt ctc gaa tca	1104
60	Ile Gly Asp Ser Pro Ile Asp Thr Phe Leu Met Glu Val Leu Glu Ser	
	355 360 365	
65	tct tca cat gac gtt caa gta gct aca	1131
	Ser Ser His Asp Val Gln Val Ala Thr	
70	370 375	
75	<210> 12	
	<211> 377	
80	<212> PRT	
	<213> Myzus persicae	
85	<400> 12	
	Met Tyr Ser Asn Ser Tyr Thr Met Tyr Ser Ser Asp Arg Leu Tyr Ser	
90	1 5 10 15	

P:\OPER\MRO\ECDY\$ONE.CIP - 1/7/99

- 31 -

	Val	Asp	Arg	Asn	Ser	Met	Met	Asn	Asn	Ser	Cys	Asn	Val	Gln	Asp	Ser
				20						25					30	
5																
	Pro	Asn	Tyr	Pro	Pro	Asn	His	Pro	Leu	Ser	Gly	Ser	Lys	His	Leu	Cys
			35					40					45			
10																
	Ser	Ile	Cys	Gly	Asp	Arg	Ala	Ser	Gly	Lys	His	Tyr	Gly	Val	Tyr	Ser
15		50					55					60				
20	Cys	Glu	Gly	Cys	Lys	Gly	Phe	Phe	Lys	Arg	Thr	Val	Arg	Lys	Asn	Leu
	65					70					75					80
25	Ser	Tyr	Ala	Cys	Arg	Glu	Glu	Asn	Lys	Cys	Ile	Ile	Asp	Lys	Arg	Gln
					85					90					95	
30	Arg	Asn	Arg	Cys	Gln	Tyr	Cys	Arg	Tyr	Gln	Lys	Cys	Leu	Thr	Met	Gly
				100					105					110		
35																
	Met	Lys	Arg	Glu	Ala	Val	Gln	Glu	Glu	Arg	Gln	Arg	Thr	Lys	Glu	Arg
40			115					120					125			
	Asp	His	Asn	Asn	Ile	Glu	Val	Glu	Pro	Thr	Ser	Ser	Ser	Asn	Thr	Asp
45		130					135					140				
50	Met	Pro	Val	Glu	Leu	Ile	Leu	Arg	Ala	Glu	Asn	Lys	Ala	Asp	Ala	Ile
	145					150					155					160
55	Lys	Thr	Glu	Gln	Gln	Tyr	Ile	Glu	Gln	Arg	His	Pro	Gln	His	Thr	Val
				165					170					175		
60	Gly	Ala	Ile	Cys	Gln	Ala	Thr	Asp	Lys	Gln	Leu	Ile	Gln	Leu	Val	Glu
				180					185					190		
65																
	Trp	Ala	Lys	His	Ile	Pro	His	Phe	Lys	Asn	Leu	Pro	Leu	Gly	Asp	Gln
70			195					200					205			
	Val	Leu	Leu	Leu	Arg	Ala	Gly	Trp	Asn	Glu	Leu	Met	Ile	Ala	Ala	Phe
75		210					215					220				
80	Ser	His	Arg	Ser	Ile	Ser	Val	Lys	Asp	Gly	Ile	Val	Leu	Ala	Thr	Gly
	225					230					235					240
85	Leu	Thr	Val	Asp	Arg	Asp	Ser	Ala	His	Gln	Ala	Gly	Val	Glu	Ala	Ile
					245					250						

- 32 -

```

260                               265                               270
5  Met Asp Arg Thr Glu Leu Gly Cys Leu Arg Thr Ile Ile Leu Phe Asn
    275                               280                               285

10 Pro Gly Ser Lys Gly Leu Gln Ser Val Asn Glu Val Gln Val Leu Arg
    290                               295                               300

15 Asp Lys Val Tyr Val Ala Leu Glu Glu Tyr Cys Arg Thr Thr His Pro
    305                               310                               315                               320

20 Glu Glu Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ser Leu
    325                               330                               335

25 Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Tyr Lys Leu
    340                               345                               350

30 Ile Gly Asp Ser Pro Ile Asp Thr Phe Leu Met Glu Val Leu Glu Ser
    355                               360                               365

40 Ser Ser His Asp Val Gln Val Ala Thr
    370                               375

45

<210> 13
<211> 150
<212> DNA
50 <213> Lucilia cuprina

<220>
<221> CDS
<222> (9)..(134)

55
<400> 13
aattctgc gaa gga tgc aag gga ttc ttc aaa cgt acc gta cgc aag gac 50
    Glu Gly Cys Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp
        1                               5                               10

60
ttg aca tat gct tgt cgt gag gac aga aat tgc att ata gat aaa cga 98

```

- 33 -

Leu Thr Tyr Ala Cys Arg Glu Asp Arg Asn Cys Ile Ile Asp Lys Arg

15

20

25

30

caa aga aat cgt tgc cag tat tgt cgc tac caa aag tgatcgatac cgtcga 150

5 Gln Arg Asn Arg Cys Gln Tyr Cys Arg Tyr Gln Lys

35

40

<210> 14

10 <211> 42

<212> PRT

<213> Lucilia cuprina

<400> 14

Glu Gly Cys Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp Leu Thr

15

1

5

10

15

Tyr Ala Cys Arg Glu Asp Arg Asn Cys Ile Ile Asp Lys Arg Gln Arg

20

25

30

20 Asn Arg Cys Gln Tyr Cys Arg Tyr Gln Lys

35

40

<210> 15

25 <211> 32

<212> DNA

<213> Artificial Sequence

<220>

30 <223> Description of Artificial Sequence:primer

ccpccccc

[illegible]

32

<213> Artificial Sequence

32

23

<212> DNA

<213> Artificial Sequence

<223> Description of Artificial Sequence:primer

<400> 18

gcactccTga cactttcgTc tca

23

10 <210> 19

<211> 23

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence:primer

<400> 19

20 t c g t c c g g t t a c c a t t a c a a c g c

23

<210> 20

<211> 24

25 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

30 <400> 20

tagacotttg geraaytcna caat

24

JOINT INVENTORS' DECLARATION FOR PATENT APPLICATION
AND POWER OF ATTORNEY

As the below named inventors, we hereby declare that:

Our residences, post office addresses and citizenship are as stated below our names.

We believe that we are the original, first and joint inventors of the subject matter which is claimed and for which a patent is sought on the invention entitled: "Genetic Sequences Encoding Steroid and Juvenile Hormone Receptor Polypeptides and Insecticidal Modalities Therefor," the specification of which was filed on 1 July 1999, as Application Serial No. _____;

We hereby authorize our legal representative to add reference to the Serial No. and/or filing date of the above-referenced application to this declaration.

We hereby state that we have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, §1.56.

Prior Foreign Application(s)

We hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application(s) for patent or inventor's certificate having a filing date before that of the application to which priority is claimed:

Country	Application No.	Date of Filing (day,month,year)	Date of Issue (day,month,year)	Priority Claimed 35 U.S.C.119
AU	PP1356/98	15 January 1999		Yes <u>X</u> No <u> </u>

Prior Provisional Application(s)

We hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below:

Application Serial Number	Date of Filing (day,month,year)
------------------------------	------------------------------------

Prior U.S. Application(s) and PCT International Application(s) Designating the United States

We hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s), or § 365(c) of any PCT International application(s) designating the United States listed below:

Application Serial Number	Date of Filing (day,month,year)	Status(Patented,Pending,Abandoned)
PCT/AU99/00033	15 January 1999	Pending

Insofar as the subject matter of each of the claims in this application is not disclosed in the prior United States, foreign or PCT International application(s) to which priority has been claimed above in the manner provided by the first paragraph of Title 35, United States Code, §112, we acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application.

We hereby appoint, both jointly and severally, as our attorneys and agents with full power of substitution and revocation, to prosecute this application and any corresponding application filed in the Patent Cooperation Treaty Receiving Office, and to transact all business in the Patent and Trademark Office connected herewith the following attorneys and agents, their registration numbers being listed after their names:

Lorance L. Greenlee, Reg. No. 27,894; Ellen P. Winner, Reg. No. 28,547; Sally A. Sullivan, Reg. No. 32,064; Donna M. Ferber, Reg. No. 33,878; No. 37,374; G. William VanCleave, Reg. No. 40,213; and Susan K. Doughty, Reg. No. 43,595, all of Greenlee,

Winner and Sullivan, P.C., 5370 Manhattan Circle, Suite 201, Boulder, CO 80303.

We hereby declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

(1) Full Name of

First Inventor: HILL, Ronald Johnston
Residence: Forestville, New South Wales, 2087, Australia
Citizenship: Australia
Post Office Address: Altona Avenue, Forestville, New South Wales, 2087, Australia

(1) Signature _____ Date _____

(2) Full Name of

Second Inventor: HANNAN, Garry Noel
Residence: Hunters Hill, New South Wales, 2110, Australia
Citizenship: Australia
Post Office Address: 32 Earl Street, Hunters Hill, New South Wales, 2110, Australia

(2) Signature _____ Date _____